



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 223/12, 211/76, 213/75, A61K 31/55, 31/445, 31/44	A1	(11) International Publication Number: WO 96/29313 (43) International Publication Date: 26 September 1996 (26.09.96)
(21) International Application Number: PCT/US96/03726 (22) International Filing Date: 19 March 1996 (19.03.96) (30) Priority Data: 08/407,839 21 March 1995 (21.03.95) US (71) Applicant: THE PROCTER & GAMBLE COMPANY [US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US). (72) Inventors: DE, Biswanath; 11269 Cornell Woods Drive, Cincinnati, OH 45241 (US). CHENG, Menyen; 5860 Hazel- tine Court, West Chester, OH 45069 (US). NATCHUS, Michael, George; 11269 Corona Road, Cincinnati, OH 45250 (US). WAHL, Christopher, Thomas; 5238 Mulhauser Road, Hamilton, OH 45011 (US). (74) Agents: REED, T., David et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45217 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>
(54) Title: LACTAM-CONTAINING HYDROXAMIC ACID DERIVATIVES, THEIR PREPARATION AND THEIR USE AS IN- HIBITORS OF MATRIX METALLOPROTEASE		
<div style="text-align: center;"> </div> <div style="text-align: right;">(I)</div>		
(57) Abstract <p>The present invention relates to compounds that exhibit inhibitory activity against matrix metalloproteases ("MMPs"). Because MMPs are known to play a role in tissue degradation, the compounds of the present invention may be useful in preventing or treating diseases associated with excess MMP activity. In particular, the compounds have a structure according to formula (I), wherein (A) (1) (a) R¹ is hydrogen; alkyl; heteroalkyl; alkenyl; a heterocyclic ring; a carbocyclic ring; alkoxy; carbocycle-alkyl; heterocycle-alkyl; carbocycle-heteroalkyl; or heterocycle-heteroalkyl; and (b) R² is hydrogen; hydroxy; alkyl; alkenyl; alkynyl; heteroalkyl; a heterocyclic ring; a carbocyclic ring; carbocycle-alkyl; heterocycle-alkyl; or -OR, where R is alkyl, alkenyl, or carbocycle-alkyl; or (2) R¹ and R² together form a cycloalkyl ring having from 3 to 8 ring atoms; (B) R³ is hydrogen; alkyl; or carbocycle-alkyl; (C) R⁴ is: (1) alkyl; (2) carbocycle-alkyl; (3) -X-C(=Y)-Z-R⁵ or -X-CH₂-Z-R⁵, where (a) X is covalent bond or alkyl; (b) Y is O, S, or NH; (c) Z is O, S, or NH; and (d) R⁵ is hydrogen; alkyl; alkenyl; carbocycle-alkyl; or aryl; or (4) -SO₂-R⁶, where R⁶ is alkyl, carbocycle-alkyl, heterocycle-alkyl, or aryl; and (D) Q is -[C(R⁷)₂]-_n, where (1) n is the integer 2, 3, or 4; and (2) each R⁷ is independently hydrogen or alkyl so the Q-containing heterocycle is saturated; or the R⁷ moiety on two adjacent carbon atoms is a covalent bond such that the Q-containing heterocycle in Formula (I) is unsaturated; or a pharmaceutically-acceptable salt; or biohydrolyzable alkoxyamide, acyloxyamide, or imide thereof. The invention also relates to pharmaceutical compositions comprising these compounds, and methods for preventing or treating diseases associated with unwanted MMP activity using the compounds and compositions.</p>		



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 223/12, 211/76, 213/75, A61K 31/55, 31/445, 31/44		A1	(11) International Publication Number: WO 96/29313
			(43) International Publication Date: 26 September 1996 (26.09.96)
(21) International Application Number: PCT/US96/03726 (22) International Filing Date: 19 March 1996 (19.03.96) (30) Priority Data: 08/407,839 21 March 1995 (21.03.95) US (71) Applicant: THE PROCTER & GAMBLE COMPANY [US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US). (72) Inventors: DE, Biswanath; 11269 Cornell Woods Drive, Cincinnati, OH 45241 (US). CHENG, Menyen; 5860 Hazeltine Court, West Chester, OH 45069 (US). NATCHUS, Michael, George; 11269 Corona Road, Cincinnati, OH 45250 (US). WAHL, Christopher, Thomas; 5238 Mulhauser Road, Hamilton, OH 45011 (US). (74) Agents: REED, T., David et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45217 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: LACTAM-CONTAINING HYDROXAMIC ACID DERIVATIVES, THEIR PREPARATION AND THEIR USE AS INHIBITORS OF MATRIX METALLOPROTEASE			
<div style="text-align: center;"> <p style="text-align: right;">(I)</p> </div>			
(57) Abstract <p>The present invention relates to compounds that exhibit inhibitory activity against matrix metalloproteases ("MMPs"). Because MMPs are known to play a role in tissue degradation, the compounds of the present invention may be useful in preventing or treating diseases associated with excess MMP activity. In particular, the compounds have a structure according to formula (I), wherein (A) (1) (a) R¹ is hydrogen; alkyl; heteroalkyl; alkenyl; a heterocyclic ring; a carbocyclic ring; alkoxy; carbocycle-alkyl; heterocycle-alkyl; carbocycle-heteroalkyl; or heterocycle-heteroalkyl; and (b) R² is hydrogen; hydroxy; alkyl; alkenyl; alkynyl; heteroalkyl; a heterocyclic ring; a carbocyclic ring; carbocycle-alkyl; heterocycle-alkyl; or -OR, where R is alkyl, alkenyl, or carbocycle-alkyl; or (2) R¹ and R² together form a cycloalkyl ring having from 3 to 8 ring atoms; (B) R³ is hydrogen; alkyl; or carbocycle-alkyl; (C) R⁴ is: (1) alkyl; (2) carbocycle-alkyl; (3) -X-C(=Y)-Z-R⁵ or -X-CH₂-Z-R⁵, where (a) X is covalent bond or alkyl; (b) Y is O, S, or NH; (c) Z is O, S, or NH; and (d) R⁵ is hydrogen; alkyl; alkenyl; carbocycle-alkyl; or aryl; or (4) -SO₂-R⁶, where R⁶ is alkyl, carbocycle-alkyl, heterocycle-alkyl, or aryl; and (D) Q is -[C(R⁷)₂]-_n, where (1) n is the integer 2, 3, or 4; and (2) each R⁷ is independently hydrogen or alkyl so the Q-containing heterocycle is saturated; or the R⁷ moiety on two adjacent carbon atoms is a covalent bond such that the Q-containing heterocycle in Formula (I) is unsaturated; or a pharmaceutically-acceptable salt; or biohydrolyzable alkoxyamide, acyloxyamide, or imide thereof. The invention also relates to pharmaceutical compositions comprising these compounds, and methods for preventing or treating diseases associated with unwanted MMP activity using the compounds and compositions.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LJ	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LJ	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

**LACTAM-CONTAINING HYDROXAMIC ACID DERIVATIVES: THEIR PREPARATION AND THEIR
USE AS INHIBITORS OF MATRIX METALLOPROTEASE**

TECHNICAL FIELD

This invention is directed to compounds which are useful in treating diseases associated with excess and/or unwanted matrix metalloprotease activity, particularly collagenase and/or stromelysin activity. More specifically, the invention is directed to hydroxamic acid compounds that contain a substituted lactam ring.

BACKGROUND

A number of enzymes effect the breakdown of structural proteins and are structurally related metalloproteases. These include human skin fibroblast collagenase, human skin fibroblast gelatinase, human sputum collagenase and gelatinase, and human stromelysin. These are zinc-containing metalloprotease enzymes, as are the angiotensin-converting enzymes and the enkephalinases. Collagenase, stromelysin and related enzymes are important in mediating the symptomatology of a number of diseases, including rheumatoid arthritis (Mullins, D. E., et al., *Biochim Biophys Acta* (1983) 695:117-214); osteoarthritis (Henderson, B., et al., *Drugs of the Future* (1990) 15:495-508); the metastasis of tumor cells (ibid, Broadhurst, M. J., et al., European Patent Application 276,436 (published 1987), Reich, R., et al., 48 Cancer Res 3307-3312 (1988); and various ulcerated conditions. Ulcerative conditions can result in the cornea as the result of alkali burns or as a result of infection by *Pseudomonas aeruginosa*, *Acanthamoeba*, Herpes simplex and vaccinia viruses.

Other conditions characterized by unwanted matrix metalloprotease activity include periodontal disease, epidermolysis bullosa and scleritis. In view of the involvement of matrix metalloproteases in a number of disease conditions, attempts have been made to prepare inhibitors to these enzymes. A number of such inhibitors are disclosed in the literature. Examples include U.S. Patent No. 5,183,900, issued February 2, 1993 to Galardy; U.S. Patent No. 4,996,358, issued February 26, 1991 to Handa, et al.; U.S. Patent No. 4,771,038, issued September 13, 1988 to Wolanin, et al.; U.S. Patent Number 4,743,587, issued May 10, 1988 to Dickens, et al., European Patent Publication Number 575,844, published December 29, 1993 by Broadhurst, et al.; International Patent Publication No. WO 93/09090, published May 13, 1993 by Isomura, et al.; World Patent Publication 92/17460, published October 15, 1992 by

LACTAM-CONTAINING HYDROXAMIC ACID DERIVATIVES; THEIR PREPARATION AND THEIR USE AS INHIBITORS OF MATRIX METALLOPROTEASE

TECHNICAL FIELD

This invention is directed to compounds which are useful in treating diseases associated with excess and/or unwanted matrix metalloprotease activity, particularly collagenase and/or stromelysin activity. More specifically, the invention is directed to hydroxamic acid compounds that contain a substituted lactam ring.

BACKGROUND

A number of enzymes effect the breakdown of structural proteins and are structurally related metalloproteases. These include human skin fibroblast collagenase, human skin fibroblast gelatinase, human sputum collagenase and gelatinase, and human stromelysin. These are zinc-containing metalloprotease enzymes, as are the angiotensin-converting enzymes and the enkephalinases. Collagenase, stromelysin and related enzymes are important in mediating the symptomatology of a number of diseases, including rheumatoid arthritis (Mullins, D. E., et al., *Biochim Biophys Acta* (1983) 695:117-214); osteoarthritis (Henderson, B., et al., *Drugs of the Future* (1990) 15:495-508); the metastasis of tumor cells (ibid, Broadhurst, M. J., et al., European Patent Application 276,436 (published 1987), Reich, R., et al., 48 Cancer Res 3307-3312 (1988); and various ulcerated conditions. Ulcerative conditions can result in the cornea as the result of alkali burns or as a result of infection by *Pseudomonas aeruginosa*, *Acanthamoeba*, Herpes simplex and vaccinia viruses.


Other conditions characterized by unwanted matrix metalloprotease activity include periodontal disease, epidermolysis bullosa and scleritis. In view of the involvement of matrix metalloproteases in a number of disease conditions, attempts have been made to prepare inhibitors to these enzymes. A number of such inhibitors are disclosed in the literature. Examples include U.S. Patent No. 5,183,900, issued February 2, 1993 to Galaray; U.S. Patent No. 4,996,358, issued February 26, 1991 to Handa, et al.; U.S. Patent No. 4,771,038, issued September 13, 1988 to Wolanin, et al.; U.S. Patent Number 4,743,587, issued May 10, 1988 to Dickens, et al., European Patent Publication Number 575,844, published December 29, 1993 by Broadhurst, et al.; International Patent Publication No. WO 93/09090, published May 13, 1993 by Isomura, et al.; World Patent Publication 92/17460, published October 15, 1992 by

Markwell et al.; and European Patent Publication Number 498,665, published August 12, 1992 by Beckett, et al.

It is well known in the art that inhibitors of matrix metalloproteases are useful in treating diseases caused, at least in part, by breakdown of structural proteins. Though a variety of inhibitors have been prepared, there is a continuing need for potent matrix metalloprotease inhibitors useful in treating such diseases. Applicants have found that, surprisingly, the lactam-containing hydroxamic acids of the present invention are potent inhibitors of collagenase and/or stromelysin. The compounds of the present invention therefore may be useful for the treatment of conditions and diseases which are characterized by unwanted activity by the class of proteins which destroy structural proteins.

SUMMARY OF THE INVENTION

The invention provides compounds which are useful as inhibitors of matrix metalloproteases, and which are effective in treating conditions characterized by excess activity of these enzymes. In particular, the present invention relates to a compound having a structure according to Formula (I)



wherein


- (A) (1) (a) R^1 is hydrogen; alkyl; heteroalkyl; alkenyl; a heterocyclic ring; a carbocyclic ring; alkoxy; carbocycle-alkyl; heterocycle-alkyl; carbocycle-heteroalkyl; or heterocycle-heteroalkyl; and
- (b) R^2 is hydrogen; hydroxy; alkyl; alkenyl; alkynyl; heteroalkyl; a heterocyclic ring; a carbocyclic ring; carbocycle-alkyl; heterocycle-alkyl; or -OR, where R is alkyl, alkenyl, or carbocycle-alkyl; or
- (2) R^1 and R^2 together form a cycloalkyl ring having from 3 to 8 ring atoms;
- (B) R^3 is hydrogen; alkyl; or carbocycle-alkyl;
- (C) R^4 is
 - (1) alkyl;

Markwell et al.; and European Patent Publication Number 498,665, published August 12, 1992 by Beckett, et al.

It is well known in the art that inhibitors of matrix metalloproteases are useful in treating diseases caused, at least in part, by breakdown of structural proteins. Though a variety of inhibitors have been prepared, there is a continuing need for potent matrix metalloprotease inhibitors useful in treating such diseases. Applicants have found that, surprisingly, the lactam-containing hydroxamic acids of the present invention are potent inhibitors of collagenase and/or stromelysin. The compounds of the present invention therefore may be useful for the treatment of conditions and diseases which are characterized by unwanted activity by the class of proteins which destroy structural proteins.

SUMMARY OF THE INVENTION

The invention provides compounds which are useful as inhibitors of matrix metalloproteases, and which are effective in treating conditions characterized by excess activity of these enzymes. In particular, the present invention relates to a compound having a structure according to Formula (I)



wherein

- (A) (1) (a) R^1 is hydrogen; alkyl; heteroalkyl; alkenyl; a heterocyclic ring; a carbocyclic ring; alkoxy; carbocycle-alkyl; heterocycle-alkyl; carbocycle-heteroalkyl; or heterocycle-heteroalkyl; and
- (b) R^2 is hydrogen; hydroxy; alkyl; alkenyl; alkynyl; heteroalkyl; a heterocyclic ring; a carbocyclic ring; carbocycle-alkyl; heterocycle-alkyl; or -OR, where R is alkyl, alkenyl, or carbocycle-alkyl; or
- (2) R^1 and R^2 together form a cycloalkyl ring having from 3 to 8 ring atoms;
- (B) R^3 is hydrogen; alkyl; or carbocycle-alkyl;
- (C) R^4 is
 - (1) alkyl;

- (2) carbocycle-alkyl;
- (3) $-X-C(=Y)-Z-R^5$ or $-X-CH_2-Z-R^5$, where
 - (a) X is covalent bond or alkyl;
 - (b) Y is O, S, or NH;
 - (c) Z is O, S, or NH; and
 - (d) R^5 is hydrogen, alkyl, alkenyl, carbocycle-alkyl, or aryl;
 or
- (4) $-SO_2-R^6$, where R^6 is alkyl, carbocycle-alkyl, heterocycle-alkyl, or aryl; and
- (D) Q is an alkyl chain, an alkenyl chain, a heteroalkyl chain, or a heteroalkenyl chain; wherein said chain has 2, 3, or 4 chain atoms and is unsubstituted or substituted with one or more alkyl moieties;

or a pharmaceutically-acceptable salt, or biohydrolyzable alkoxyamide, acyloxyamide, or imide thereof.

These compounds have the ability to inhibit at least one mammalian matrix metalloprotease. Accordingly, in other aspects, the invention is directed to pharmaceutical compositions containing the compounds of Formula (I), and to methods of treating diseases characterized by matrix metalloprotease activity using these compounds or the pharmaceutical compositions containing them.

Matrix metalloproteases at a particularly undesired location can be targeted by conjugating the compounds of the invention to a targeting ligand specific for a marker at that location such as an antibody or fragment thereof or a receptor ligand.

The invention is also directed to various other processes which take advantage of the unique properties of these compounds. Thus, in another aspect, the invention is directed to the compounds of Formula (I) conjugated to solid supports. These conjugates can be used as affinity reagents for the purification of a desired matrix metalloprotease.

In another aspect, the invention is directed to the compounds of Formula (I) conjugated to label. As the compounds of the invention bind to at least one matrix metalloprotease, the label can be used to detect the presence of relatively high levels of matrix metalloprotease in vivo or in vitro cell culture.

In addition, the compounds of Formula (I) can be conjugated to carriers which permit the use of these compounds in immunization protocols to prepare antibodies specifically immunoreactive with the compounds of the invention. These antibodies are then useful both in therapy and in monitoring the dosage of the inhibitors.

DETAILED DESCRIPTION

- (2) carbocycle-alkyl;
 - (3) $-X-C(=Y)-Z-R^5$ or $-X-CH_2-Z-R^5$, where
 - (a) X is covalent bond or alkyl;
 - (b) Y is O, S, or NH;
 - (c) Z is O, S, or NH; and
 - (d) R^5 is hydrogen; alkyl; alkenyl; carbocycle-alkyl; or aryl;
 or
 - (4) $-SO_2-R^6$, where R^6 is alkyl, carbocycle-alkyl, heterocycle-alkyl, or aryl; and
 - (D) Q is an alkyl chain, an alkenyl chain, a heteroalkyl chain, or a heteroalkenyl chain; wherein said chain has 2, 3, or 4 chain atoms and is unsubstituted or substituted with one or more alkyl moieties;
- or a pharmaceutically-acceptable salt, or biohydrolyzable alkoxyamide, acyloxyamide, or imide thereof.

These compounds have the ability to inhibit at least one mammalian matrix metalloprotease. Accordingly, in other aspects, the invention is directed to pharmaceutical compositions containing the compounds of Formula (I), and to methods of treating diseases characterized by matrix metalloprotease activity using these compounds or the pharmaceutical compositions containing them.

Matrix metalloproteases at a particularly undesired location can be targeted by conjugating the compounds of the invention to a targeting ligand specific for a marker at that location such as an antibody or fragment thereof or a receptor ligand.


The invention is also directed to various other processes which take advantage of the unique properties of these compounds. Thus, in another aspect, the invention is directed to the compounds of Formula (I) conjugated to solid supports. These conjugates can be used as affinity reagents for the purification of a desired matrix metalloprotease.

In another aspect, the invention is directed to the compounds of Formula (I) conjugated to label. As the compounds of the invention bind to at least one matrix metalloprotease, the label can be used to detect the presence of relatively high levels of matrix metalloprotease in vivo or in vitro cell culture.

In addition, the compounds of Formula (I) can be conjugated to carriers which permit the use of these compounds in immunization protocols to prepare antibodies specifically immunoreactive with the compounds of the invention. These antibodies are then useful both in therapy and in monitoring the dosage of the inhibitors.

DETAILED DESCRIPTION


The compounds of the present invention are inhibitors of mammalian matrix metalloproteases. Preferably, the compounds are those of Formula (I) where the Q-containing heterocycle has one nitrogen atom. These compounds are the following



wherein

- (A) (1) (a) R¹ is hydrogen; alkyl; heteroalkyl; alkenyl; a heterocyclic ring; a carbocyclic ring; alkoxy; carbocycle-alkyl; heterocycle-alkyl; carbocycle-heteroalkyl; or heterocycle-heteroalkyl; and
- (b) R² is hydrogen; hydroxy; alkyl; alkenyl; alkynyl; heteroalkyl; a heterocyclic ring; a carbocyclic ring; carbocycle-alkyl; heterocycle-alkyl; or -OR, where R is alkyl, alkenyl, or carbocycle-alkyl; or
- (2) R¹ and R² together form a cycloalkyl ring having from 3 to 8 ring atoms;
- (B) R³ is hydrogen; alkyl; or carbocycle-alkyl;
- (C) R⁴ is
 - (1) alkyl;
 - (2) carbocycle-alkyl;
 - (3) -X-C(=Y)-Z-R⁵ or -X-CH₂-Z-R⁵, where
 - (a) X is covalent bond or alkyl;
 - (b) Y is O, S, or NH;
 - (c) Z is O, S, or NH; and
 - (d) R⁵ is hydrogen; alkyl; alkenyl; carbocycle-alkyl; or aryl;
 or
 - (4) -SO₂-R⁶, where R⁶ is alkyl, carbocycle-alkyl, heterocycle-alkyl, or aryl; and
- (D) Q is -[C(R⁷)₂]-_n, where
 - (1) n is the integer 2, 3, or 4; and
 - (2) each R⁷ is independently hydrogen or alkyl so the Q-containing heterocycle is saturated; or the R⁷ moiety on two adjacent carbon

The compounds of the present invention are inhibitors of mammalian matrix metalloproteases. Preferably, the compounds are those of Formula (I) where the Q-containing heterocycle has one nitrogen atom. These compounds are the following



wherein

- (A) (1) (a) R¹ is hydrogen; alkyl; heteroalkyl; alkenyl; a heterocyclic ring; a carbocyclic ring; alkoxy; carbocycle-alkyl; heterocycle-alkyl; carbocycle-heteroalkyl; or heterocycle-heteroalkyl; and
 - (b) R² is hydrogen; hydroxy; alkyl; alkenyl; alkynyl; heteroalkyl; a heterocyclic ring; a carbocyclic ring; carbocycle-alkyl; heterocycle-alkyl; or -OR, where R is alkyl, alkenyl, or carbocycle-alkyl; or
- (2) R¹ and R² together form a cycloalkyl ring having from 3 to 8 ring atoms;
- (B) R³ is hydrogen; alkyl; or carbocycle-alkyl;
- (C) R⁴ is
 - (1) alkyl;
 - (2) carbocycle-alkyl;
 - (3) -X-C(=Y)-Z-R⁵ or -X-CH₂-Z-R⁵, where
 - (a) X is covalent bond or alkyl;
 - (b) Y is O, S, or NH;
 - (c) Z is O, S, or NH; and
 - (d) R⁵ is hydrogen; alkyl; alkenyl; carbocycle-alkyl; or aryl;
 or
 - (4) -SO₂-R⁶, where R⁶ is alkyl, carbocycle-alkyl, heterocycle-alkyl, or aryl; and
- (D) Q is -[C(R⁷)₂]-_n, where
 - (1) n is the integer 2, 3, or 4; and
 - (2) each R⁷ is independently hydrogen or alkyl so the Q-containing heterocycle is saturated; or the R⁷ moiety on two adjacent carbon

atoms is a covalent bond such that the Q-containing heterocycle in Formula (I) is unsaturated;

or a pharmaceutically-acceptable salt, or biohydrolyzable alkoxyamide, acyloxyamide, or imide thereof.

Definitions and Usage of Terms:

The following is a list of definitions for terms used herein.

"Acyl" or "carbonyl" is a radical formed by removal of the hydroxy from a carboxylic acid (i.e., $R-C(=O)-$). Preferred acyl groups include (for example) acetyl, formyl, and propionyl.

"Acyloxy" is an oxygen radical having an acyl substituent (i.e., $-O$ -acyl); for example, $-O-C(=O)$ -alkyl.

"Acylamino" is an amino radical having an acyl substituent (i.e., $-N$ -acyl); for example, $-NH-C(=O)$ -alkyl.

"Alkoxyacyl" is an acyl radical ($-C(=O)-$) having an alkoxy substituent (i.e., $-O-R$), for example, $-C(=O)-O$ -alkyl.

"Alkenyl" is an unsubstituted or substituted hydrocarbon chain radical having 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 carbon atoms; preferably from 2 to 10 carbon atoms; more preferably from 2 to 8; except where indicated. Alkenyl substituents have at least one olefinic double bond (including, for example, vinyl, allyl and butenyl).

"Alkynyl" is an unsubstituted or substituted hydrocarbon chain radical having 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 carbon atoms; preferably from 2 to 10 carbon atoms; more preferably from 2 to 8; except where indicated. The chain has at least one carbon-carbon triple bond.

"Alkoxy" is an oxygen radical having a hydrocarbon chain substituent, where the hydrocarbon chain is an alkyl or alkenyl (i.e., $-O$ -alkyl or $-O$ -alkenyl). Preferred alkoxy groups include (for example) methoxy, ethoxy, propoxy and allyloxy.

"Alkoxyalkyl" is an unsubstituted or substituted alkyl moiety substituted with an alkoxy moiety (i.e., $-alkyl-O-alkyl$). Preferred is where the alkyl has 1, 2, 3, 4, 5 or 6 carbon atoms (more preferably 1 to 3 carbon atoms), and the alkoxy has 1, 2, 3, 4, 5 or 6 carbon atoms (more preferably 1 to 3 carbon atoms).

"Alkyl" is an unsubstituted or substituted saturated hydrocarbon chain radical having 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 carbon atoms; preferably from 1 to 10 carbon atoms; more preferably 1 to 4; except where indicated. Preferred alkyl groups include (for example) substituted or unsubstituted methyl, ethyl, propyl, isopropyl, and butyl.

atoms is a covalent bond such that the Q-containing heterocycle in Formula (I) is unsaturated;

or a pharmaceutically-acceptable salt, or biohydrolyzable alkoxyamide, acyloxyamide, or imide thereof.

Definitions and Usage of Terms:

The following is a list of definitions for terms used herein.

"Acyl" or "carbonyl" is a radical formed by removal of the hydroxy from a carboxylic acid (i.e., $R-C(=O)-$). Preferred acyl groups include (for example) acetyl, formyl, and propionyl.

"Acyloxy" is an oxygen radical having an acyl substituent (i.e., $-O$ -acyl); for example, $-O-C(=O)$ -alkyl.

"Acylamino" is an amino radical having an acyl substituent (i.e., $-N$ -acyl); for example, $-NH-C(=O)$ -alkyl.

"Alkoxyacyl" is an acyl radical ($-C(=O)-$) having an alkoxy substituent (i.e., $-O-R$), for example, $-C(=O)-O$ -alkyl.

"Alkenyl" is an unsubstituted or substituted hydrocarbon chain radical having 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 carbon atoms; preferably from 2 to 10 carbon atoms; more preferably from 2 to 8; except where indicated. Alkenyl substituents have at least one olefinic double bond (including, for example, vinyl, allyl and butenyl).

"Alkynyl" is an unsubstituted or substituted hydrocarbon chain radical having 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 carbon atoms; preferably from 2 to 10 carbon atoms; more preferably from 2 to 8; except where indicated. The chain has at least one carbon-carbon triple bond.

"Alkoxy" is an oxygen radical having a hydrocarbon chain substituent, where the hydrocarbon chain is an alkyl or alkenyl (i.e., $-O$ -alkyl or $-O$ -alkenyl). Preferred alkoxy groups include (for example) methoxy, ethoxy, propoxy and allyloxy.

"Alkoxyalkyl" is an unsubstituted or substituted alkyl moiety substituted with an alkoxy moiety (i.e., $-alkyl-O-alkyl$). Preferred is where the alkyl has 1, 2, 3, 4, 5 or 6 carbon atoms (more preferably 1 to 3 carbon atoms), and the alkoxy has 1, 2, 3, 4, 5 or 6 carbon atoms (more preferably 1 to 3 carbon atoms).

"Alkyl" is an unsubstituted or substituted saturated hydrocarbon chain radical having 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 carbon atoms; preferably from 1 to 10 carbon atoms; more preferably 1 to 4; except where indicated. Preferred alkyl groups include (for example) substituted or unsubstituted methyl, ethyl, propyl, isopropyl, and butyl.

"Alkylamino" is an amino radical having one (secondary amine) or two (tertiary amine) alkyl substituents (i.e., -N-alkyl). For example, methylamine (-NHCH₃), dimethylamine (-N(CH₃)₂), methylethylamine (-N(CH₃)CH₂CH₃).

"Aminoacyl" is acyl radical having an amino substituent (i.e., -C(=O)-N); for example, -C(=O)-NH₂. The amino group of the aminoacyl moiety may be unsubstituted (i.e., primary amine) or may be substituted with one (secondary amine) or two (i.e., tertiary amine) alkyl groups.

"Aryl" is an aromatic carbocyclic ring radical. Preferred aryl groups include (for example) phenyl, tolyl, xylyl, cumenyl and naphthyl.

"Arylalkyl" is an alkyl radical substituted with an aryl group. Preferred arylalkyl groups include benzyl, phenylethyl, and phenylpropyl.

"Arylalkylamino" is an amine radical substituted with an arylalkyl group (e.g., -NH-benzyl).

"Arylamino" is an amine radical substituted with an aryl group (i.e., -NH-aryl).

"Aryloxy" is an oxygen radical having an aryl substituent (i.e., -O-aryl).

"Carbocyclic ring" is an unsubstituted or substituted, saturated, unsaturated or aromatic, hydrocarbon ring radical. Carbocyclic rings are monocyclic or are fused, bridged or spiro polycyclic ring systems. Monocyclic carbocyclic rings generally contain 3, 4, 5, 6, 7, 8 or 9 atoms, preferably 3 to 6 atoms. Polycyclic carbocyclic rings contain 7, 8, 9, 10, 11, 12, 13, 14, 15, 16 or 17 atoms, preferably from 7 to 13 atoms.

"Carbocycle-alkyl" is an unsubstituted or substituted alkyl radical substituted with a carbocyclic ring. Unless otherwise specified, the carbocyclic ring is preferably an aryl or cycloalkyl; more preferably an aryl. Preferred carbocycle-alkyl groups include benzyl, phenylethyl and phenylpropyl.

"Carbocycle-heteroalkyl" is an unsubstituted or substituted heteroalkyl radical substituted with a carbocyclic ring. Unless otherwise specified, the carbocyclic ring is preferably an aryl or cycloalkyl; more preferably an aryl. The heteroalkyl is preferably 2-oxa-propyl, 2-oxa-ethyl, 2-thia-propyl, or 2-thia-ethyl.

"Carboxyalkyl" is an unsubstituted or substituted alkyl radical substituted with a carboxy (-C(=O)OH) moiety. For example, -CH₂-C(=O)OH.

"Cycloalkyl" is a saturated carbocyclic ring radical. Preferred cycloalkyl groups include (for example) cyclopropyl, cyclobutyl and cyclohexyl.

"Cycloheteroalkyl" is a saturated heterocyclic ring. Preferred cycloheteroalkyl groups include (for example) morpholine, piperidine, and piperazine.

"Alkylamino" is an amino radical having one (secondary amine) or two (tertiary amine) alkyl substituents (i.e., -N-alkyl). For example, methylamine (-NHCH₃), dimethylamine (-N(CH₃)₂), methylethylamine (-N(CH₃)CH₂CH₃).

"Aminoacyl" is acyl radical having an amino substituent (i.e., -C(=O)-N); for example, -C(=O)-NH₂. The amino group of the aminoacyl moiety may be unsubstituted (i.e., primary amine) or may be substituted with one (secondary amine) or two (i.e., tertiary amine) alkyl groups.

"Aryl" is an aromatic carbocyclic ring radical. Preferred aryl groups include (for example) phenyl, tolyl, xylyl, cumenyl and naphthyl.

"Arylalkyl" is an alkyl radical substituted with an aryl group. Preferred arylalkyl groups include benzyl, phenylethyl, and phenylpropyl.

"Arylalkylamino" is an amine radical substituted with an arylalkyl group (e.g., -NH-benzyl).

"Arylamino" is an amine radical substituted with an aryl group (i.e., -NH-aryl).

"Aryloxy" is an oxygen radical having an aryl substituent (i.e., -O-aryl).

"Carbocyclic ring" is an unsubstituted or substituted, saturated, unsaturated or aromatic, hydrocarbon ring radical. Carbocyclic rings are monocyclic or are fused, bridged or spiro polycyclic ring systems. Monocyclic carbocyclic rings generally contain 3, 4, 5, 6, 7, 8 or 9 atoms, preferably 3 to 6 atoms. Polycyclic carbocyclic rings contain 7, 8, 9, 10, 11, 12, 13, 14, 15, 16 or 17 atoms, preferably from 7 to 13 atoms.

"Carbocycle-alkyl" is an unsubstituted or substituted alkyl radical substituted with a carbocyclic ring. Unless otherwise specified, the carbocyclic ring is preferably an aryl or cycloalkyl; more preferably an aryl. Preferred carbocycle-alkyl groups include benzyl, phenylethyl and phenylpropyl.

"Carbocycle-heteroalkyl" is an unsubstituted or substituted heteroalkyl radical substituted with a carbocyclic ring. Unless otherwise specified, the carbocyclic ring is preferably an aryl or cycloalkyl; more preferably an aryl. The heteroalkyl is preferably 2-oxa-propyl, 2-oxa-ethyl, 2-thia-propyl, or 2-thia-ethyl.

"Carboxyalkyl" is an unsubstituted or substituted alkyl radical substituted with a carboxy (-C(=O)OH) moiety. For example, -CH₂-C(=O)OH.

"Cycloalkyl" is a saturated carbocyclic ring radical. Preferred cycloalkyl groups include (for example) cyclopropyl, cyclobutyl and cyclohexyl.

"Cycloheteroalkyl" is a saturated heterocyclic ring. Preferred cycloheteroalkyl groups include (for example) morpholine, piperadine, and piperazine.

"Fused rings" are rings that are superimposed together such that they share two ring atoms. A given ring may be fused to more than one other ring.

"Heterocycle-alkyl" is an alkyl radical substituted with a heterocyclic ring. The heterocyclic ring is preferably an heteroaryl or cycloheteroalkyl; more preferably an heteroaryl.

"Heterocycle-heteroalkyl" is an unsubstituted or substituted heteroalkyl radical substituted with a heterocyclic ring. The heterocyclic ring is preferably an aryl or cycloheteroalkyl; more preferably an aryl.

"Heteroatom" is a nitrogen, sulfur or oxygen atom. Groups containing one or more heteroatoms may contain different heteroatoms.

"Heteroalkenyl" is an unsubstituted or substituted unsaturated chain radical having 3, 4, 5, 6, 7 or 8 members comprising carbon atoms and one or two heteroatoms. The chain has at least one carbon-carbon double bond.

"Heteroalkyl" is an unsubstituted or substituted saturated chain radical having 2, 3, 4, 5, 6, 7 or 8 comprising carbon atoms and one or two heteroatoms.

"Heterocyclic ring" is an unsubstituted or substituted, saturated, unsaturated or aromatic ring radical comprised of carbon atoms and one or more heteroatoms in the ring. Heterocyclic rings are monocyclic or are fused, bridged or spiro polycyclic ring systems. Monocyclic heterocyclic rings contain 3, 4, 5, 6, 7, 8 or 9 atoms, preferably 4 to 7 atoms. Polycyclic rings contain 7, 8, 9, 10, 11, 12, 13, 14, 15, 16 or 17 atoms, preferably from 7 to 13 atoms.

"Heteroaryl" is an aromatic heterocyclic ring radical. Preferred heteroaryl groups include (for example) thienyl, furyl, pyrrolyl, pyridinyl, pyrazinyl, thiazolyl, pyrimidinyl, quinolinyl, and tetrazolyl.

"Halo", "halogen", or "halide" is a chloro, bromo, fluoro or iodo atom radical. Chloro and fluoro are preferred halides.

Also, as referred to herein, a "lower" hydrocarbon moiety (e.g., "lower" alkyl) is a hydrocarbon chain comprised of 1, 2, 3, 4, 5 or 6, preferably from 1 to 4, carbon atoms.

A "pharmaceutically-acceptable salt" is a cationic salt formed at any acidic (e.g., carboxyl) group, or an anionic salt formed at any basic (e.g., amino) group. Many such salts are known in the art, as described in World Patent Publication 87/05297, Johnston et al., published September 11, 1987 (incorporated by reference herein). Preferred cationic salts include the alkali metal salts (such as sodium and potassium), and alkaline earth metal salts (such as magnesium and calcium). Preferred anionic salts include the halides (such as chloride salts).

"Fused rings" are rings that are superimposed together such that they share two ring atoms. A given ring may be fused to more than one other ring.

"Heterocycle-alkyl" is an alkyl radical substituted with a heterocyclic ring. The heterocyclic ring is preferably an heteroaryl or cycloheteroalkyl; more preferably an heteroaryl.

"Heterocycle-heteroalkyl" is an unsubstituted or substituted heteroalkyl radical substituted with a heterocyclic ring. The heterocyclic ring is preferably an aryl or cycloheteroalkyl; more preferably an aryl.

"Heteroatom" is a nitrogen, sulfur or oxygen atom. Groups containing one or more heteroatoms may contain different heteroatoms.

"Heteroalkenyl" is an unsubstituted or substituted unsaturated chain radical having 3, 4, 5, 6, 7 or 8 members comprising carbon atoms and one or two heteroatoms. The chain has at least one carbon-carbon double bond.

"Heteroalkyl" is an unsubstituted or substituted saturated chain radical having 2, 3, 4, 5, 6, 7 or 8 comprising carbon atoms and one or two heteroatoms.

"Heterocyclic ring" is an unsubstituted or substituted, saturated, unsaturated or aromatic ring radical comprised of carbon atoms and one or more heteroatoms in the ring. Heterocyclic rings are monocyclic or are fused, bridged or spiro polycyclic ring systems. Monocyclic heterocyclic rings contain 3, 4, 5, 6, 7, 8 or 9 atoms, preferably 4 to 7 atoms. Polycyclic rings contain 7, 8, 9, 10, 11, 12, 13, 14, 15, 16 or 17 atoms, preferably from 7 to 13 atoms.


"Heteroaryl" is an aromatic heterocyclic ring radical. Preferred heteroaryl groups include (for example) thienyl, furyl, pyrrolyl, pyridinyl, pyrazinyl, thiazolyl, pyrimidinyl, quinolinyl, and tetrazolyl.

"Halo", "halogen", or "halide" is a chloro, bromo, fluoro or iodo atom radical. Chloro and fluoro are preferred halides.

Also, as referred to herein, a "lower" hydrocarbon moiety (e.g., "lower" alkyl) is a hydrocarbon chain comprised of 1, 2, 3, 4, 5 or 6, preferably from 1 to 4, carbon atoms.


A "pharmaceutically-acceptable salt" is a cationic salt formed at any acidic (e.g., carboxyl) group, or an anionic salt formed at any basic (e.g., amino) group. Many such salts are known in the art, as described in World Patent Publication 87/05297, Johnston et al., published September 11, 1987 (incorporated by reference herein). Preferred cationic salts include the alkali metal salts (such as sodium and potassium), and alkaline earth metal salts (such as magnesium and calcium). Preferred anionic salts include the halides (such as chloride salts).

"Biohydrolyzable alkoxyamide" and "Biohydrolyzable acyloxyamide" are amides of a hydroxamic acid that do not essentially interfere with the inhibitory activity of the compound, or that are readily converted in vivo by a human or lower animal subject to yield an active hydroxamic acid. A biohydrolyzable alkoxyamide derivative of the Formula (I) compounds is represented by the following:



where E is an alkyl moiety. A biohydrolyzable acyloxyamide derivative of the Formula (I) compounds is where E is an acyl moiety (e.g. $R-C(=O)-$).

A "biohydrolyzable hydroxy imide" is an imide of a Formula (I) compound that does not interfere with the metalloprotease inhibitory activity of these compounds, or that is readily converted in vivo by a human or lower animal subject to yield an active Formula (I) compound. Such hydroxy imides include those that do not interfere with the biological activity of the Formula (I) compounds. These imides have a structure according to the following:




where E is an acyl moiety (e.g., -C(=O)-R).

A "solvate" is a complex formed by the combination of a solute (e.g., a hydroxamic acid) and a solvent (e.g., water). See J. Honig et al., The Van Nostrand Chemist's Dictionary, p. 650 (1953). Pharmaceutically-acceptable solvents used according to this invention include those that do not interfere with the biological activity of the hydroxamic acid (e.g., water, ethanol, acetic acid, N,N-dimethylformamide).

The illustration of specific protected forms and other derivatives of the Formula (I) compounds is not intended to be limiting. The application of other useful protecting groups, salt forms, etc. is within the ability of the skilled artisan.


As defined above and as used herein, substituent groups may themselves be substituted. Such substitution may be with one or more substituents. Such substituents include those listed in C. Hansch and A. Leo, Substituent Constants for

"Biohydrolyzable alkoxyamide" and "Biohydrolyzable acyloxyamide" are amides of a hydroxamic acid that do not essentially interfere with the inhibitory activity of the compound, or that are readily converted in vivo by a human or lower animal subject to yield an active hydroxamic acid. A biohydrolyzable alkoxyamide derivative of the Formula (I) compounds is represented by the following:



where E is an alkyl moiety. A biohydrolyzable acyloxyamide derivative of the Formula (I) compounds is where E is an acyl moiety (e.g. $R-C(=O)-$).

A "biohydrolyzable hydroxy imide" is an imide of a Formula (I) compound that does not interfere with the metalloprotease inhibitory activity of these compounds, or that is readily converted in vivo by a human or lower animal subject to yield an active Formula (I) compound. Such hydroxy imides include those that do not interfere with the biological activity of the Formula (I) compounds. These imides have a structure according to the following:



where E is an acyl moiety (e.g., -C(=O)-R).

A "solvate" is a complex formed by the combination of a solute (e.g., a hydroxamic acid) and a solvent (e.g., water). See J. Honig et al., The Van Nostrand Chemist's Dictionary, p. 650 (1953). Pharmaceutically-acceptable solvents used according to this invention include those that do not interfere with the biological activity of the hydroxamic acid (e.g., water, ethanol, acetic acid, N,N-dimethylformamide).

The illustration of specific protected forms and other derivatives of the Formula (I) compounds is not intended to be limiting. The application of other useful protecting groups, salt forms, etc. is within the ability of the skilled artisan.

As defined above and as used herein, substituent groups may themselves be substituted. Such substitution may be with one or more substituents. Such substituents include those listed in C. Hansch and A. Leo, Substituent Constants for

Correlation Analysis in Chemistry and Biology (1979), incorporated by reference herein. Preferred substituents include (for example) alkyl, alkenyl, alkoxy, hydroxy, oxo, nitro, amino, aminoalkyl (e.g., aminomethyl, etc.), cyano, halo, carboxy, alkoxyaceyl (e.g., carboethoxy, etc.), thiol, aryl, cycloalkyl, heteroaryl, heterocycloalkyl (e.g., piperidiny, morpholinyl, pyrrolidinyl, etc.), imino, thioxo, hydroxyalkyl, aryloxy, arylalkyl, and combinations thereof.

As used herein, "mammalian matrix metalloprotease" means any metal-containing enzyme found in mammalian sources which is capable of catalyzing the breakdown of collagen, gelatin or proteoglycan under suitable assay conditions. Appropriate assay conditions can be found, for example, in U.S. Pat. No. 4,743,587, which references the procedure of Cawston, et al., *Anal Biochem* (1979) 99:340-345, use of a synthetic substrate is described by Weingarten, H., et al., *Biochem Biophys Res Comm* (1984) 139:1184-1187. Any standard method for analyzing the breakdown of these structural proteins can, of course, be used. The matrix metalloprotease enzymes referred to herein are all zinc-containing proteases which are similar in structure to, for example, human stromelysin or skin fibroblast collagenase. The ability of candidate compounds to inhibit matrix metalloprotease activity can, of course, be tested in the assays described above. Isolated matrix metalloprotease enzymes can be used to confirm the inhibiting activity of the invention compounds, or crude extracts which contain the range of enzymes capable of tissue breakdown can be used.

Compounds:

Referring to Formula (I), the R^1 substituent group is selected from hydrogen; alkyl; heteroalkyl; alkenyl; a heterocyclic ring; a carbocyclic ring; alkoxy; carbocycle-alkyl; heterocycle-alkyl; carbocycle-heteroalkyl; and heterocycle-heteroalkyl. Preferred is where R^1 is hydrogen; alkyl; alkenyl; a heterocyclic ring; alkoxy; carbocycle-alkyl; or aminoalkyl. More preferred is where R^1 is hydrogen; C_1 - C_8 alkyl; aminoalkyl; or benzyl. Most preferred is where R^1 is hydrogen, methyl, ethyl, or propyl.

R^2 is selected from hydrogen; hydroxy; alkyl; alkenyl; alkynyl; heteroalkyl; a heterocyclic ring; a carbocyclic ring; carbocycle-alkyl; heterocycle-alkyl; and -OR, where R is alkyl, alkenyl, or carbocycle-alkyl. Preferred is where R^2 is hydrogen; alkyl; or aminoalkyl. More preferred is where R^2 is hydrogen or C_1 - C_8 alkyl. Particularly preferred is where R^2 is n-octyl, n-pentyl or 2-methylpropyl.

In the alternative, R^1 and R^2 can together form a cycloalkyl ring having from 3 to 8 ring atoms; preferably 5 to 7 ring atoms; more preferably 6 atoms. Preferred is where R^1 and R^2 do not combine to form a ring.

Correlation Analysis in Chemistry and Biology (1979), incorporated by reference herein. Preferred substituents include (for example) alkyl, alkenyl, alkoxy, hydroxy, oxo, nitro, amino, aminoalkyl (e.g., aminomethyl, etc.), cyano, halo, carboxy, alkoxyaceyl (e.g., carboethoxy, etc.), thiol, aryl, cycloalkyl, heteroaryl, heterocycloalkyl (e.g., piperidiny, morpholinyl, pyrrolidinyl, etc.), imino, thioxo, hydroxyalkyl, aryloxy, arylalkyl, and combinations thereof.

As used herein, "mammalian matrix metalloprotease" means any metal-containing enzyme found in mammalian sources which is capable of catalyzing the breakdown of collagen, gelatin or proteoglycan under suitable assay conditions. Appropriate assay conditions can be found, for example, in U.S. Pat. No. 4,743,587, which references the procedure of Cawston, et al., *Anal Biochem* (1979) 99:340-345, use of a synthetic substrate is described by Weingarten, H., et al., *Biochem Biophys Res Comm* (1984) 139:1184-1187. Any standard method for analyzing the breakdown of these structural proteins can, of course, be used. The matrix metalloprotease enzymes referred to herein are all zinc-containing proteases which are similar in structure to, for example, human stromelysin or skin fibroblast collagenase. The ability of candidate compounds to inhibit matrix metalloprotease activity can, of course, be tested in the assays described above. Isolated matrix metalloprotease enzymes can be used to confirm the inhibiting activity of the invention compounds, or crude extracts which contain the range of enzymes capable of tissue breakdown can be used.

Compounds:

Referring to Formula (I), the R^1 substituent group is selected from hydrogen; alkyl; heteroalkyl; alkenyl; a heterocyclic ring; a carbocyclic ring; alkoxy; carbocycle-alkyl; heterocycle-alkyl; carbocycle-heteroalkyl; and heterocycle-heteroalkyl. Preferred is where R^1 is hydrogen; alkyl; alkenyl; a heterocyclic ring; alkoxy; carbocycle-alkyl; or aminoalkyl. More preferred is where R^1 is hydrogen; C_1 - C_8 alkyl; aminoalkyl; or benzyl. Most preferred is where R^1 is hydrogen, methyl, ethyl, or propyl.

R^2 is selected from hydrogen; hydroxy; alkyl; alkenyl; alkynyl; heteroalkyl; a heterocyclic ring; a carbocyclic ring; carbocycle-alkyl; heterocycle-alkyl; and -OR, where R is alkyl, alkenyl, or carbocycle-alkyl. Preferred is where R^2 is hydrogen; alkyl; or aminoalkyl. More preferred is where R^2 is hydrogen or C_1 - C_8 alkyl. Particularly preferred is where R^2 is n-octyl, n-pentyl or 2-methylpropyl.

In the alternative, R^1 and R^2 can together form a cycloalkyl ring having from 3 to 8 ring atoms; preferably 5 to 7 ring atoms; more preferably 6 atoms. Preferred is where R^1 and R^2 do not combine to form a ring.

R^3 is selected from hydrogen; alkyl; and carbocycle-alkyl (more preferably C_1 - C_2 alkyl). Preferred is where R^3 is hydrogen.

R^4 is selected from alkyl; carbocycle-alkyl; alkoxyalkyl; $-X-C(=Y)-Z-R^5$ or $-X-CH_2-Z-R^5$, where (a) X is covalent bond or alkyl; (b) Y is O, S, or NH; (c) Z is O, S, or NH; R^5 is hydrogen; alkyl, alkenyl, carbocycle-alkyl, or aryl.

When R^4 is alkyl, preferred is C_1 - C_8 alkyl.


When R^4 is $-X-C(=Y)-Z-R^5$, X is preferably C_1 - C_3 alkyl (more preferably C_1 - C_2 alkyl), Y is preferably O, and Z is preferably NH or O. When Y and Z are both O, R^5 is preferably alkyl (preferably methyl or ethyl; most preferably methyl) or carbocycle-alkyl (preferably benzyl); most preferably alkyl. When Y is O and Z is NH, R^5 is preferably alkyl or carbocycle-alkyl; more preferably methyl, ethyl, butyl, or benzyl.

When R^4 is $-X-CH_2-Z-R^5$, X is preferably C_1 - C_3 alkyl, Z is preferably O or S, and R^5 is preferably alkyl or carbocycle-alkyl (more preferably alkyl). Particularly preferred is where X is C_1 , Z is O and R^5 is C_1 - C_3 alkyl.

When R^4 is $-SO_2R^6$, R^6 is alkyl, carbocycle-alkyl, heterocycle-alkyl, or aryl; preferably aryl (preferably phenyl; most preferably 4-methylphenyl).

As indicated above, particularly preferred compounds of the present invention are those where the Q-containing heterocycle has only one ring nitrogen atom. That is, where Q is $[-C(R^7)_2-]_n$, where n is the integer 2, 3, or 4 (more preferably 3 or 4). Particularly preferred is where n is 4, such that the Q-containing heterocycle has 7 ring atoms. Each R^7 is independently hydrogen or alkyl; or the R^7 moiety on two adjacent carbon atoms is a covalent bond such that the Q-containing heterocycle in Formula (I) is unsaturated. Preferred compounds are those where the Q-containing heterocycle is saturated; most preferably where each R^7 is hydrogen.

The following illustrates compounds where the Q-containing heterocycle is unsaturated:



In this structure, the heterocycle has seven members (i.e., $n=4$). Referring to Formula (I), carbon atoms *a* and *b* each represent a $-C(R^7)_2-$ moiety where one R^7 is hydrogen and the other is a covalent bond, such that a double bond exists between atoms *a* and *b*.

Two groups of adjacent carbon atoms may have R^7 moieties that are covalent bonds, such that the lactam ring has two points of unsaturation (i.e., two double bonds).

R^3 is selected from hydrogen; alkyl; and carbocycle-alkyl (more preferably C_1 - C_2 alkyl). Preferred is where R^3 is hydrogen.

R^4 is selected from alkyl; carbocycle-alkyl; alkoxyalkyl; $-X-C(=Y)-Z-R^5$ or $-X-CH_2-Z-R^5$, where (a) X is covalent bond or alkyl; (b) Y is O, S, or NH; (c) Z is O, S, or NH; R^5 is hydrogen; alkyl, alkenyl, carbocycle-alkyl, or aryl.

When R^4 is alkyl, preferred is C_1 - C_8 alkyl.


When R^4 is $-X-C(=Y)-Z-R^5$, X is preferably C_1 - C_3 alkyl (more preferably C_1 - C_2 alkyl), Y is preferably O, and Z is preferably NH or O. When Y and Z are both O, R^5 is preferably alkyl (preferably methyl or ethyl; most preferably methyl) or carbocycle-alkyl (preferably benzyl); most preferably alkyl. When Y is O and Z is NH, R^5 is preferably alkyl or carbocycle-alkyl; more preferably methyl, ethyl, butyl, or benzyl.

When R^4 is $-X-CH_2-Z-R^5$, X is preferably C_1 - C_3 alkyl, Z is preferably O or S, and R^5 is preferably alkyl or carbocycle-alkyl (more preferably alkyl). Particularly preferred is where X is C_1 , Z is O and R^5 is C_1 - C_3 alkyl.

When R^4 is $-SO_2R^6$, R^6 is alkyl, carbocycle-alkyl, heterocycle-alkyl, or aryl; preferably aryl (preferably phenyl; most preferably 4-methylphenyl).

As indicated above, particularly preferred compounds of the present invention are those where the Q-containing heterocycle has only one ring nitrogen atom. That is, where Q is $[-C(R^7)_2-]_n$, where n is the integer 2, 3, or 4 (more preferably 3 or 4). Particularly preferred is where n is 4, such that the Q-containing heterocycle has 7 ring atoms. Each R^7 is independently hydrogen or alkyl; or the R^7 moiety on two adjacent carbon atoms is a covalent bond such that the Q-containing heterocycle in Formula (I) is unsaturated. Preferred compounds are those where the Q-containing heterocycle is saturated; most preferably where each R^7 is hydrogen.


The following illustrates compounds where the Q-containing heterocycle is unsaturated:



In this structure, the heterocycle has seven members (i.e., $n=4$). Referring to Formula (I), carbon atoms *a* and *b* each represent a $-C(R^7)_2-$ moiety where one R^7 is hydrogen and the other is a covalent bond, such that a double bond exists between atoms *a* and *b*.

Two groups of adjacent carbon atoms may have R^7 moieties that are covalent bonds, such that the lactam ring has two points of unsaturation (i.e., two double bonds).

The following illustrates such rings where the Q-containing heterocycle has two points of unsaturation:




In this structure, the heterocycle has six members (i.e., $n = 3$). Referring to Formula (I), carbon atoms a and b each represent a $-C(R^7)_2-$ moiety where one R^7 is hydrogen and the other is a covalent bond, such that a double bond exists between atoms a and b . In addition, c and d each represents a $-C(R^7)_2-$ moiety where one R^7 is hydrogen and the other is a covalent bond, such that a double bond exists between atoms c and d .

The following table lists representative preferred compounds within the scope of the invention. The table is not intended to be an exhaustive list of the compounds within the scope of the invention. Referring to Formula (I), Q is $(-CH_2-)_n$, n is 4, and R^3 is hydrogen in each instance.

Cmp.	R^1	R^2	R^4
1	hydrogen	2-methylpropyl	$-CH_2-C(=O)-O-CH_3$
2	hydrogen	2-methylpropyl	$-CH_2-C(=O)-NH-CH_3$
3	hydrogen	2-methylpropyl	$-CH_2-C(=O)-O-C(CH_3)_3$
4	hydrogen	2-methylpropyl	$-CH_2-phenyl$
5	hydrogen	2-methylpropyl	$-CH_2-C(=O)-O-CH_2-phenyl$
6	hydrogen	2-methylpropyl	$-CH_2-C(=O)-NH-CH_2-phenyl$
7	hydrogen	2-methylpropyl	$-CH_2-C(=O)-O-CH_3$
8	hydrogen	$-(CH_2)_4CH_3$	$-CH_2-C(=O)-NH-(CH_2)_3-CH_3$
9	hydrogen	$-(CH_2)_7CH_3$	$-CH_2-C(=O)-O-CH_3$
10	hydrogen	$-(CH_2)_7CH_3$	$-SO_2-phenyl$
11	hydrogen	$-(CH_2)_7CH_3$	$-CH_2-CH_2-O-CH_3$
12	hydrogen	$-(CH_2)_7CH_3$	$-(CH_2)_3CH_3$
13	CH_3 (S form)	2-methylpropyl	$-CH_2-C(=O)-O-CH_3$
14	CH_3 (R form)	2-methylpropyl	$-CH_2-C(=O)-O-CH_3$
15	$CH_3CH_2CH_2-$	2-methylpropyl	$-CH_2-C(=O)-O-CH_3$
16	$-(CH_2)_2-CH_2OH$ (S form)	2-methylpropyl	$-CH_2-C(=O)-O-CH_3$
17	CH_3 (S form)	$-(CH_2)_7CH_3$	$-CH_2-C(=O)-O-CH_3$
18	CH_3 (R form)	$-(CH_2)_7CH_3$	$-CH_2-C(=O)-O-CH_3$
19	CH_3 (S form)	$-(CH_2)_7CH_3$	$-CH_2-CH_2-O-CH_3$

The following illustrates such rings where the Q-containing heterocycle has two points of unsaturation:






In this structure, the heterocycle has six members (i.e., $n = 3$). Referring to Formula (I), carbon atoms *a* and *b* each represent a $-C(R^7)_2-$ moiety where one R^7 is hydrogen and the other is a covalent bond, such that a double bond exists between atoms *a* and *b*. In addition, *c* and *d* each represents a $-C(R^7)_2-$ moiety where one R^7 is hydrogen and the other is a covalent bond, such that a double bond exists between atoms *c* and *d*.

The following table lists representative preferred compounds within the scope of the invention. The table is not intended to be an exhaustive list of the compounds within the scope of the invention. Referring to Formula (I), Q is $(-CH_2-)_n$, n is 4, and R^3 is hydrogen in each instance.

Cmp.	R^1	R^2	R^4
1	hydrogen	2-methylpropyl	$-CH_2-C(=O)-O-CH_3$
2	hydrogen	2-methylpropyl	$-CH_2-C(=O)-NH-CH_3$
3	hydrogen	2-methylpropyl	$-CH_2-C(=O)-O-C(CH_3)_3$
4	hydrogen	2-methylpropyl	$-CH_2$ -phenyl
5	hydrogen	2-methylpropyl	$-CH_2-C(=O)-O-CH_2$ -phenyl
6	hydrogen	2-methylpropyl	$-CH_2-C(=O)-NH-CH_2$ -phenyl
7	hydrogen	2-methylpropyl	$-CH_2-C(=O)-O-CH_3$
8	hydrogen	$-(CH_2)_4CH_3$	$-CH_2-C(=O)-NH-(CH_2)_3-CH_3$
9	hydrogen	$-(CH_2)_7CH_3$	$-CH_2-C(=O)-O-CH_3$
10	hydrogen	$-(CH_2)_7CH_3$	$-SO_2$ -phenyl
11	hydrogen	$-(CH_2)_7CH_3$	$-CH_2-CH_2-O-CH_3$
12	hydrogen	$-(CH_2)_7CH_3$	$-(CH_2)_3CH_3$
13	CH_3 (S form)	2-methylpropyl	$-CH_2-C(=O)-O-CH_3$
14	CH_3 (R form)	2-methylpropyl	$-CH_2-C(=O)-O-CH_3$
15	$CH_3CH_2CH_2-$	2-methylpropyl	$-CH_2-C(=O)-O-CH_3$
16	$-(CH_2)_2-CH_2OH$ (S form)	2-methylpropyl	$-CH_2-C(=O)-O-CH_3$
17	CH_3 (S form)	$-(CH_2)_7CH_3$	$-CH_2-C(=O)-O-CH_3$
18	CH_3 (R form)	$-(CH_2)_7CH_3$	$-CH_2-C(=O)-O-CH_3$
19	CH_3 (S form)	$-(CH_2)_7CH_3$	$-CH_2-CH_2-O-CH_3$




20 CH₃ (R form)-(CH₂)₇CH₃-CH₂-CH₂-O-CH₃General Schemes for Compound Preparation:

The hydroxamic compounds of Formula (I) can be prepared using a variety of procedures. General schemes include the following. (Representative examples are described for making specific compounds hereinbelow.)

a. General Scheme 1:1) (BOC)₂O, DMSO; 2A) LiN(TMS)₂; R⁴-X, THF;2B) t-BuOK, R⁴-X, DMF; 3) TFA/CH₂Cl₂ or HCl/Et₂O4) LiN(TMS)₂, THF, R¹-X; 5) LDA, THF6) EDAC, HOBT, NMM, DMF, 0 deg. C; 7) TFA, CH₂Cl₂;8A-i) EDAC, HOBT, NMM, DMF, BnONH₂·HCl, 0 deg. C; ii) H₂/Pd-C, EtOH;8B-i) CH₂N₂; ii) NH₂OH·HCl/KOH, MeOH

20 CH₃ (R form)-(CH₂)₇CH₃-CH₂-CH₂-O-CH₃General Schemes for Compound Preparation:

The hydroxamic compounds of Formula (I) can be prepared using a variety of procedures. General schemes include the following. (Representative examples are described for making specific compounds hereinbelow.)


a. General Scheme 1:1) (BOC)₂O, DMSO; 2A) LiN(TMS)₂; R⁴-X, THF;2B) t-BuOK, R⁴-X, DMF; 3) TFA/CH₂Cl₂ or HCl/Et₂O4) LiN(TMS)₂, THF, R¹-X; 5) LDA, THF6) EDAC, HOBT, NMM, DMF, 0 deg. C; 7) TFA, CH₂Cl₂;8A-i) EDAC, HOBT, NMM, DMF, BnONH₂·HCl, 0 deg. C; ii) H₂/Pd-C, EtOH;8B-i) CH₂N₂; ii) NH₂OH·HCl/KOH, MeOH

Commercially available Caprolactam (A) is protected to give (B), followed by alkylation of the amide nitrogen under appropriate conditions to give (C). The derivatized lactam (C) is deprotected under acidic conditions to give the amine salt (D) which is then used for coupling to various succinates as described in Scheme 2 and 3.

The various alkyl succinates (E) are synthesized following Evan's chiral alkylation method (D. A. Evans, et al., *Org. Synth.* Vol. 86, p 83 (1990), incorporated herein by reference). The dianion generated by treating (E) with a hindered base is alkylated to give syn-disubstituted succinates (F) which on further treatment with LDA gives the desired anti-diastereomer (G) in reasonable yield (H. J. Crimmin, et al., *Synlett*, 137-138 (1993)).

The acid (F) or (G) and the amine salt (D) are coupled under a mild condition to give the amide (H) (depicted without specifying stereochemistry), which on deprotection under acidic conditions gives the corresponding acid (I). A final transformation is carried out to convert the purified acid to the desired inhibitor (J).

b. General Scheme 2:




A direct alkylation method can also be utilized to synthesize the final inhibitors. For example, the treatment of intermediate (K) (prepared by reacting Compound A with Compound F or G according to Scheme 1) with a hindered base at low temperature followed by a quench of the anion with an alkylating agent gives (L) which on deprotection under acidic conditions and re-esterification provides (M) in good yield. A direct treatment of this ester with freshly-generated hydroxylamine then produces the final inhibitor (N).

Commercially available Caprolactam (A) is protected to give (B), followed by alkylation of the amide nitrogen under appropriate conditions to give (C). The derivatized lactam (C) is deprotected under acidic conditions to give the amine salt (D) which is then used for coupling to various succinates as described in Scheme 2 and 3.

The various alkyl succinates (E) are synthesized following Evan's chiral alkylation method (D. A. Evans, et al., *Org. Synth.* Vol. 86, p 83 (1990), incorporated herein by reference). The dianion generated by treating (E) with a hindered base is alkylated to give syn-disubstituted succinates (F) which on further treatment with LDA gives the desired anti-diastereomer (G) in reasonable yield (H. J. Crimmin, et al., *Synlett*, 137-138 (1993)).

The acid (F) or (G) and the amine salt (D) are coupled under a mild condition to give the amide (H) (depicted without specifying stereochemistry), which on deprotection under acidic conditions gives the corresponding acid (I). A final transformation is carried out to convert the purified acid to the desired inhibitor (J).


b. General Scheme 2:



A direct alkylation method can also be utilized to synthesize the final inhibitors. For example, the treatment of intermediate (K) (prepared by reacting Compound A with Compound F or G according to Scheme 1) with a hindered base at low temperature followed by a quench of the anion with an alkylating agent gives (L) which on deprotection under acidic conditions and re-esterification provides (M) in good yield. A direct treatment of this ester with freshly-generated hydroxylamine then produces the final inhibitor (N).

c. General Scheme 3:

The compounds of the present invention having a lactam ring with 5 or 6 members can be prepared as follows.




- 1) $(\text{TMS})_2\text{NH}$, CH_3CN ; 2) $(\text{BOC})_2\text{O}$; DMSO
 3) $\text{LiN}(\text{TMS})_2$, THF, $\text{BrCH}_2\text{COOMe}$; 4) TFA, CH_2Cl_2

For example, L-ornithine hydrochloride (O) on heating under reflux provides the six-membered lactam (P) which on further protection with BOC-anhydride produces the desired amide (Q) in reasonable yield. This intermediate can then be carried on to the final product (S) following a method as described in the preceding schemes. For 5-membered lactams, L-ornithine hydrochloride (Compound (O)) is replaced by $(\text{COOH})\text{CH}(\text{NH}_2)\text{CH}_2\text{CH}_2\text{NH}_2$ as the starting material.

d. General Scheme 4:


Modifications of the ring system described in General Scheme 3 can also be made via the following a method, to provide unsaturation in the Q-containing ring.



Here, a properly-substituted pyrimidone (T) is N-alkylated under appropriate conditions to give (U). The resultant product, in this case the nitro-pyrimidone (U), is reduced to provide the desired amine (V). This intermediate is coupled to the succinate (F) or (G) described in Scheme 1 (see Scheme 1 for the synthesis of final products). A variety of ring systems can be generated in a similar fashion.

c. General Scheme 3:

The compounds of the present invention having a lactam ring with 5 or 6 members can be prepared as follows.




- 1) $(\text{TMS})_2\text{NH}$, CH_3CN ; 2) $(\text{BOC})_2\text{O}$; DMSO
 3) $\text{LiN}(\text{TMS})_2$, THF, $\text{BrCH}_2\text{COOMe}$; 4) TFA, CH_2Cl_2

For example, L-ornithine hydrochloride (O) on heating under reflux provides the six-membered lactam (P) which on further protection with BOC-anhydride produces the desired amide (Q) in reasonable yield. This intermediate can then be carried on to the final product (S) following a method as described in the preceding schemes. For 5-membered lactams, L-ornithine hydrochloride (Compound (O)) is replaced by $(\text{COOH})\text{CH}(\text{NH}_2)\text{CH}_2\text{CH}_2\text{NH}_2$ as the starting material.

d. General Scheme 4:

Modifications of the ring system described in General Scheme 3 can also be made via the following a method, to provide unsaturation in the Q-containing ring.



Here, a properly-substituted pyrimidone (T) is N-alkylated under appropriate conditions to give (U). The resultant product, in this case the nitro-pyrimidone (U), is reduced to provide the desired amine (V). This intermediate is coupled to the succinate (F) or (G) described in Scheme 1 (see Scheme 1 for the synthesis of final products). A variety of ring systems can be generated in a similar fashion.

Compositions:

The compositions of the invention comprise:

- (a) a safe and effective amount of a compound of Formula (I); and
- (b) a pharmaceutically-acceptable carrier.

As discussed above, numerous diseases are known to be mediated by excess or undesired matrix-destroying metalloprotease activity. These include tumor metastasis, osteoarthritis, rheumatoid arthritis, skin inflammation, ulcerations, particularly of the cornea, reaction to infection, periodontitis and the like. Thus, the compounds of the invention are useful in therapy with regard to conditions involving this unwanted activity.

The invention compounds can therefore be formulated into pharmaceutical compositions for use in treatment or prophylaxis of these conditions. Standard pharmaceutical formulation techniques are used, such as those disclosed in Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, Pa., latest edition.

A "safe and effective amount" of a Formula (I) compound is an amount that is effective, to inhibit matrix metalloproteases at the site(s) of activity, in a human or lower animal subject, without undue adverse side effects (such as toxicity, irritation, or allergic response), commensurate with a reasonable benefit/risk ratio when used in the manner of this invention. The specific "safe and effective amount" will, obviously, vary with such factors as the particular condition being treated, the physical condition of the patient, the duration of treatment, the nature of concurrent therapy (if any), the specific dosage form to be used, the carrier employed, the solubility of the Formula (I) compound therein, and the dosage regimen desired for the composition.

The compositions of this invention are preferably provided in unit dosage form. As used herein, a "unit dosage form" is a composition of this invention containing an amount of a Formula (I) compound that is suitable for administration to a human or lower animal subject, in a single dose, according to good medical practice. These compositions preferably contain from about 5 mg (milligrams) to about 1000 mg, more preferably from about 10 mg to about 500 mg, more preferably from about 10 mg to about 300 mg, of a Formula (I) compound.

The compositions of this invention may be in any of a variety of forms, suitable (for example) for oral, rectal, topical or parenteral administration. Depending upon the particular route of administration desired, a variety of pharmaceutically-acceptable carriers well-known in the art may be used. These include solid or liquid fillers, diluents, hydrotropes, surface-active agents, and

Compositions:

The compositions of the invention comprise:

- (a) a safe and effective amount of a compound of Formula (I); and
- (b) a pharmaceutically-acceptable carrier.

As discussed above, numerous diseases are known to be mediated by excess or undesired matrix-destroying metalloprotease activity. These include tumor metastasis, osteoarthritis, rheumatoid arthritis, skin inflammation, ulcerations, particularly of the cornea, reaction to infection, periodontitis and the like. Thus, the compounds of the invention are useful in therapy with regard to conditions involving this unwanted activity.

The invention compounds can therefore be formulated into pharmaceutical compositions for use in treatment or prophylaxis of these conditions. Standard pharmaceutical formulation techniques are used, such as those disclosed in Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, Pa., latest edition.

A "safe and effective amount" of a Formula (I) compound is an amount that is effective, to inhibit matrix metalloproteases at the site(s) of activity, in a human or lower animal subject, without undue adverse side effects (such as toxicity, irritation, or allergic response), commensurate with a reasonable benefit/risk ratio when used in the manner of this invention. The specific "safe and effective amount" will, obviously, vary with such factors as the particular condition being treated, the physical condition of the patient, the duration of treatment, the nature of concurrent therapy (if any), the specific dosage form to be used, the carrier employed, the solubility of the Formula (I) compound therein, and the dosage regimen desired for the composition.

The compositions of this invention are preferably provided in unit dosage form. As used herein, a "unit dosage form" is a composition of this invention containing an amount of a Formula (I) compound that is suitable for administration to a human or lower animal subject, in a single dose, according to good medical practice. These compositions preferably contain from about 5 mg (milligrams) to about 1000 mg, more preferably from about 10 mg to about 500 mg, more preferably from about 10 mg to about 300 mg, of a Formula (I) compound.

The compositions of this invention may be in any of a variety of forms, suitable (for example) for oral, rectal, topical or parenteral administration. Depending upon the particular route of administration desired, a variety of pharmaceutically-acceptable carriers well-known in the art may be used. These include solid or liquid fillers, diluents, hydrotropes, surface-active agents, and

encapsulating substances. Optional pharmaceutically-active materials may be included, which do not substantially interfere with the inhibitory activity of the Formula (I) compound. The amount of carrier employed in conjunction with the Formula (I) compound is sufficient to provide a practical quantity of material for administration per unit dose of the Formula (I) compound. Techniques and compositions for making dosage forms useful in the methods of this invention are described in the following references, all incorporated by reference herein: Modern Pharmaceutics, Chapters 9 and 10 (Banker & Rhodes, editors, 1979); Lieberman et al., Pharmaceutical Dosage Forms: Tablets (1981); and Ansel, Introduction to Pharmaceutical Dosage Forms 2d Edition (1976).

In particular, pharmaceutically-acceptable carriers for systemic administration include sugars, starches, cellulose and its derivatives, malt, gelatin, talc, calcium sulfate, vegetable oils, synthetic oils, polyols, alginic acid, phosphate buffer solutions, emulsifiers, isotonic saline, and pyrogen-free water. Preferred carriers for parenteral administration include propylene glycol, ethyl oleate, pyrrolidone, ethanol, and sesame oil. Preferably, the pharmaceutically-acceptable carrier, in compositions for parenteral administration, comprises at least about 90% by weight of the total composition.

Various oral dosage forms can be used, including such solid forms as tablets, capsules, granules and bulk powders. These oral forms comprise a safe and effective amount, usually at least about 5%, and preferably from about 25% to about 50%, of the Formula (I) compound. Tablets can be compressed, tablet triturates, enteric-coated, sugar-coated, film-coated, or multiple-compressed, containing suitable binders, lubricants, diluents, disintegrating agents, coloring agents, flavoring agents, flow-inducing agents, and melting agents. Liquid oral dosage forms include aqueous solutions, emulsions, suspensions, solutions and/or suspensions reconstituted from non-effervescent granules, and effervescent preparations reconstituted from effervescent granules, containing suitable solvents, preservatives, emulsifying agents, suspending agents, diluents, sweeteners, melting agents, coloring agents and flavoring agents.

The compositions of this invention can also be administered topically to a subject, i.e., by the direct laying on or spreading of the composition on the epidermal or epithelial tissue of the subject. Such compositions include, for example, lotions, creams, solutions, gels and solids. These topical compositions preferably comprise a safe and effective amount, usually at least about 0.1%, and preferably from about 1% to about 5%, of the Formula (I) compound. Suitable carriers for topical administration preferably remain in place on the skin as a

encapsulating substances. Optional pharmaceutically-active materials may be included, which do not substantially interfere with the inhibitory activity of the Formula (I) compound. The amount of carrier employed in conjunction with the Formula (I) compound is sufficient to provide a practical quantity of material for administration per unit dose of the Formula (I) compound. Techniques and compositions for making dosage forms useful in the methods of this invention are described in the following references, all incorporated by reference herein: Modern Pharmaceutics, Chapters 9 and 10 (Banker & Rhodes, editors, 1979); Lieberman et al., Pharmaceutical Dosage Forms: Tablets (1981); and Ansel, Introduction to Pharmaceutical Dosage Forms 2d Edition (1976).

In particular, pharmaceutically-acceptable carriers for systemic administration include sugars, starches, cellulose and its derivatives, malt, gelatin, talc, calcium sulfate, vegetable oils, synthetic oils, polyols, alginic acid, phosphate buffer solutions, emulsifiers, isotonic saline, and pyrogen-free water. Preferred carriers for parenteral administration include propylene glycol, ethyl oleate, pyrrolidone, ethanol, and sesame oil. Preferably, the pharmaceutically-acceptable carrier, in compositions for parenteral administration, comprises at least about 90% by weight of the total composition.

Various oral dosage forms can be used, including such solid forms as tablets, capsules, granules and bulk powders. These oral forms comprise a safe and effective amount, usually at least about 5%, and preferably from about 25% to about 50%, of the Formula (I) compound. Tablets can be compressed, tablet triturates, enteric-coated, sugar-coated, film-coated, or multiple-compressed, containing suitable binders, lubricants, diluents, disintegrating agents, coloring agents, flavoring agents, flow-inducing agents, and melting agents. Liquid oral dosage forms include aqueous solutions, emulsions, suspensions, solutions and/or suspensions reconstituted from non-effervescent granules, and effervescent preparations reconstituted from effervescent granules, containing suitable solvents, preservatives, emulsifying agents, suspending agents, diluents, sweeteners, melting agents, coloring agents and flavoring agents.

The compositions of this invention can also be administered topically to a subject, i.e., by the direct laying on or spreading of the composition on the epidermal or epithelial tissue of the subject. Such compositions include, for example, lotions, creams, solutions, gels and solids. These topical compositions preferably comprise a safe and effective amount, usually at least about 0.1%, and preferably from about 1% to about 5%, of the Formula (I) compound. Suitable carriers for topical administration preferably remain in place on the skin as a

continuous film, and resist being removed by perspiration or immersion in water. Generally, the carrier is organic in nature and capable of having dispersed or dissolved therein the Formula (I) compound. The carrier may include pharmaceutically-acceptable emollients, emulsifiers, thickening agents, and solvents.

Methods of Administration:

This invention also provides methods of treating or preventing disorders associated with excess or undesired matrix metalloprotease activity in a human or other animal subject, by administering a safe and effective amount of a Formula (I) compound to said subject. As used herein, a "disorder associated with excess or undesired matrix metalloprotease activity" is any disorder characterized by degradation of matrix proteins. The methods of the invention are useful in treating disorders such as (for example) osteoarthritis, periodontitis, corneal ulceration, tumor invasion, and rheumatoid arthritis.

The Formula (I) compounds and compositions of this invention can be administered topically or systemically. Systemic application includes any method of introducing Formula (I) compound into the tissues of the body, e.g., intra-articular (especially in treatment of rheumatoid arthritis), intrathecal, epidural, intramuscular, transdermal, intravenous, intraperitoneal, subcutaneous, sublingual, rectal, and oral administration. The Formula (I) compounds of the present invention are preferably administered orally.

The specific dosage of inhibitor to be administered, as well as the duration of treatment, are mutually dependent. The dosage and treatment regimen will also depend upon such factors as the specific Formula (I) compound used, the treatment indication, the ability of the Formula (I) compound to reach minimum inhibitory concentrations at the site of the matrix metalloprotease to be inhibited, the personal attributes of the subject (such as weight), compliance with the treatment regimen, and the presence and severity of any side effects of the treatment.

Typically, for a human adult (weighing approximately 70 kilograms), from about 5 mg to about 3000 mg, more preferably from about 5 mg to about 1000 mg, more preferably from about 10 mg to about 100 mg, of Formula (I) compound are administered per day. It is understood that these dosage ranges are by way of example only, and that daily administration can be adjusted depending on the factors listed above.

A preferred method of administration for treatment of rheumatoid arthritis is oral or parenterally via intra-articular injection. As is known and practiced in the art, all formulations for parenteral administration must be sterile. For mammals,

continuous film, and resist being removed by perspiration or immersion in water. Generally, the carrier is organic in nature and capable of having dispersed or dissolved therein the Formula (I) compound. The carrier may include pharmaceutically-acceptable emollients, emulsifiers, thickening agents, and solvents.

Methods of Administration:

This invention also provides methods of treating or preventing disorders associated with excess or undesired matrix metalloprotease activity in a human or other animal subject, by administering a safe and effective amount of a Formula (I) compound to said subject. As used herein, a "disorder associated with excess or undesired matrix metalloprotease activity" is any disorder characterized by degradation of matrix proteins. The methods of the invention are useful in treating disorders such as (for example) osteoarthritis, periodontitis, corneal ulceration, tumor invasion, and rheumatoid arthritis.

The Formula (I) compounds and compositions of this invention can be administered topically or systemically. Systemic application includes any method of introducing Formula (I) compound into the tissues of the body, e.g., intra-articular (especially in treatment of rheumatoid arthritis), intrathecal, epidural, intramuscular, transdermal, intravenous, intraperitoneal, subcutaneous, sublingual, rectal, and oral administration. The Formula (I) compounds of the present invention are preferably administered orally.

The specific dosage of inhibitor to be administered, as well as the duration of treatment, are mutually dependent. The dosage and treatment regimen will also depend upon such factors as the specific Formula (I) compound used, the treatment indication, the ability of the Formula (I) compound to reach minimum inhibitory concentrations at the site of the matrix metalloprotease to be inhibited, the personal attributes of the subject (such as weight), compliance with the treatment regimen, and the presence and severity of any side effects of the treatment.

Typically, for a human adult (weighing approximately 70 kilograms), from about 5 mg to about 3000 mg, more preferably from about 5 mg to about 1000 mg, more preferably from about 10 mg to about 100 mg, of Formula (I) compound are administered per day. It is understood that these dosage ranges are by way of example only, and that daily administration can be adjusted depending on the factors listed above.

A preferred method of administration for treatment of rheumatoid arthritis is oral or parenterally via intra-articular injection. As is known and practiced in the art, all formulations for parenteral administration must be sterile. For mammals,

especially humans, (assuming an approximate body weight of 70 kilograms) individual doses of from about 10 mg to about 1000 mg are preferred.

A preferred method of systemic administration is oral. Individual doses of from about 10 mg to about 1000 mg, preferably from about 10 mg to about 300 mg are preferred.

Topical administration can be used to deliver the Formula (I) compound systemically, or to treat a subject locally. The amounts of Formula (I) compound to be topically administered depends upon such factors as skin sensitivity, type and location of the tissue to be treated, the composition and carrier (if any) to be administered, the particular Formula (I) compound to be administered, as well as the particular disorder to be treated and the extent to which systemic (as distinguished from local) effects are desired.

The inhibitors of the invention can be targeted to specific locations where the matrix metalloprotease is accumulated by using targeting ligands. For example, to focus the inhibitors to matrix metalloprotease contained in a tumor, the inhibitor is conjugated to an antibody or fragment thereof which is immunoreactive with a tumor marker as is generally understood in the preparation of immunotoxins in general. The targeting ligand can also be a ligand suitable for a receptor which is present on the tumor. Any targeting ligand which specifically reacts with a marker for the intended target tissue can be used. Methods for coupling the invention compound to the targeting ligand are well known and are similar to those described below for coupling to carrier. The conjugates are formulated and administered as described above.

For localized conditions, topical administration is preferred. For example, to treat ulcerated cornea, direct application to the affected eye may employ a formulation as eyedrops or aerosol. For corneal treatment, the compounds of the invention can also be formulated as gels or ointments, or can be incorporated into collagen or a hydrophilic polymer shield. The materials can also be inserted as a contact lens or reservoir or as a subconjunctival formulation. For treatment of skin inflammation, the compound is applied locally and topically, in a gel, paste, salve or ointment. The mode of treatment thus reflects the nature of the condition and suitable formulations for any selected route are available in the art.

In all of the foregoing, of course, the compounds of the invention can be administered alone or as mixtures, and the compositions may further include additional drugs or excipients as appropriate for the indication.

Some of the compounds of the invention also inhibit bacterial metalloproteases although generally at a lower level than that exhibited with respect to mammalian metalloproteases. Some bacterial metalloproteases seem to be less dependent on the

especially humans, (assuming an approximate body weight of 70 kilograms) individual doses of from about 10 mg to about 1000 mg are preferred.

A preferred method of systemic administration is oral. Individual doses of from about 10 mg to about 1000 mg, preferably from about 10 mg to about 300 mg are preferred.

Topical administration can be used to deliver the Formula (I) compound systemically, or to treat a subject locally. The amounts of Formula (I) compound to be topically administered depends upon such factors as skin sensitivity, type and location of the tissue to be treated, the composition and carrier (if any) to be administered, the particular Formula (I) compound to be administered, as well as the particular disorder to be treated and the extent to which systemic (as distinguished from local) effects are desired.

The inhibitors of the invention can be targeted to specific locations where the matrix metalloprotease is accumulated by using targeting ligands. For example, to focus the inhibitors to matrix metalloprotease contained in a tumor, the inhibitor is conjugated to an antibody or fragment thereof which is immunoreactive with a tumor marker as is generally understood in the preparation of immunotoxins in general. The targeting ligand can also be a ligand suitable for a receptor which is present on the tumor. Any targeting ligand which specifically reacts with a marker for the intended target tissue can be used. Methods for coupling the invention compound to the targeting ligand are well known and are similar to those described below for coupling to carrier. The conjugates are formulated and administered as described above.

For localized conditions, topical administration is preferred. For example, to treat ulcerated cornea, direct application to the affected eye may employ a formulation as eyedrops or aerosol. For corneal treatment, the compounds of the invention can also be formulated as gels or ointments, or can be incorporated into collagen or a hydrophilic polymer shield. The materials can also be inserted as a contact lens or reservoir or as a subconjunctival formulation. For treatment of skin inflammation, the compound is applied locally and topically, in a gel, paste, salve or ointment. The mode of treatment thus reflects the nature of the condition and suitable formulations for any selected route are available in the art.

In all of the foregoing, of course, the compounds of the invention can be administered alone or as mixtures, and the compositions may further include additional drugs or excipients as appropriate for the indication.

Some of the compounds of the invention also inhibit bacterial metalloproteases although generally at a lower level than that exhibited with respect to mammalian metalloproteases. Some bacterial metalloproteases seem to be less dependent on the

stereochemistry of the inhibitor, whereas substantial differences are found between diastereomers in their ability to inactivate the mammalian proteases. Thus, this pattern of activity can be used to distinguish between the mammalian and bacterial enzymes.

Preparation and Use of Antibodies:

The invention compounds can also be utilized in immunization protocols to obtain antisera immunospecific for the invention compounds. As the invention compounds are relatively small, they are advantageously coupled to antigenically neutral carriers such as the conventionally used keyhole limpet hemocyanin (KLH) or serum albumin carriers. For those invention compounds having a carboxyl functionality, coupling to carrier can be done by methods generally known in the art. For example, the carboxyl residue can be reduced to an aldehyde and coupled to carrier through reaction with sidechain amino groups in protein-based carriers, optionally followed by reduction of imino linkage formed. The carboxyl residue can also be reacted with sidechain amino groups using condensing agents such as dicyclohexyl carbodiimide or other carbodiimide dehydrating agents.

Linker compounds can also be used to effect the coupling; both homobifunctional and heterobifunctional linkers are available from Pierce Chemical Company, Rockford, Ill. The resulting immunogenic complex can then be injected into suitable mammalian subjects such as mice, rabbits, and the like. Suitable protocols involve repeated injection of the immunogen in the presence of adjuvants according to a schedule which boosts production of antibodies in the serum. The titers of the immune serum can readily be measured using immunoassay procedures, now standard in the art, employing the invention compounds as antigens.

The antisera obtained can be used directly or monoclonal antibodies may be obtained by harvesting the peripheral blood lymphocytes or the spleen of the immunized animal and immortalizing the antibody-producing cells, followed by identifying the suitable antibody producers using standard immunoassay techniques.

The polyclonal or monoclonal preparations are then useful in monitoring therapy or prophylaxis regimens involving the compounds of the invention. Suitable samples such as those derived from blood, serum, urine, or saliva can be tested for the presence of the administered inhibitor at various times during the treatment protocol using standard immunoassay techniques which employ the antibody preparations of the invention.

The invention compounds can also be coupled to labels such as scintigraphic labels, e.g., technetium 99 or I-131, using standard coupling methods. The labeled compounds are administered to subjects to determine the locations of excess amounts of

stereochemistry of the inhibitor, whereas substantial differences are found between diastereomers in their ability to inactivate the mammalian proteases. Thus, this pattern of activity can be used to distinguish between the mammalian and bacterial enzymes.

Preparation and Use of Antibodies:

The invention compounds can also be utilized in immunization protocols to obtain antisera immunospecific for the invention compounds. As the invention compounds are relatively small, they are advantageously coupled to antigenically neutral carriers such as the conventionally used keyhole limpet hemocyanin (KLH) or serum albumin carriers. For those invention compounds having a carboxyl functionality, coupling to carrier can be done by methods generally known in the art. For example, the carboxyl residue can be reduced to an aldehyde and coupled to carrier through reaction with sidechain amino groups in protein-based carriers, optionally followed by reduction of imino linkage formed. The carboxyl residue can also be reacted with sidechain amino groups using condensing agents such as dicyclohexyl carbodiimide or other carbodiimide dehydrating agents.

Linker compounds can also be used to effect the coupling; both homobifunctional and heterobifunctional linkers are available from Pierce Chemical Company, Rockford, Ill. The resulting immunogenic complex can then be injected into suitable mammalian subjects such as mice, rabbits, and the like. Suitable protocols involve repeated injection of the immunogen in the presence of adjuvants according to a schedule which boosts production of antibodies in the serum. The titers of the immune serum can readily be measured using immunoassay procedures, now standard in the art, employing the invention compounds as antigens.

The antisera obtained can be used directly or monoclonal antibodies may be obtained by harvesting the peripheral blood lymphocytes or the spleen of the immunized animal and immortalizing the antibody-producing cells, followed by identifying the suitable antibody producers using standard immunoassay techniques.

The polyclonal or monoclonal preparations are then useful in monitoring therapy or prophylaxis regimens involving the compounds of the invention. Suitable samples such as those derived from blood, serum, urine, or saliva can be tested for the presence of the administered inhibitor at various times during the treatment protocol using standard immunoassay techniques which employ the antibody preparations of the invention.


The invention compounds can also be coupled to labels such as scintigraphic labels, e.g., technetium 99 or I-131, using standard coupling methods. The labeled compounds are administered to subjects to determine the locations of excess amounts of

one or more matrix metalloproteases in vivo. The ability of the inhibitors to selectively bind matrix metalloprotease is thus taken advantage of to map the distribution of these enzymes in situ. The techniques can also be employed in histological procedures and the labeled invention compounds can be used in competitive immunoassays.

The following non-limiting examples illustrate the compounds, compositions, and uses of the present invention.

Example 1

Synthesis of (2R)-Isobutyl-3-(N-hydroxycarboxamido)-propanoic acid amide of 1N-(carbomethoxymethyl)-caprolactam-(3S)-amine (1)




(2R)-Isobutyl-3-carbo-*tert*-butoxypropionic acid amide of caprolactam-(3S)-amine (25). A mixture of acid 23 (2.0 g, 8.70 mmole), caprolactam-[3S]-amine 24 (1.23 g, 9.57 mmole) and 1-Hydroxybenzotriazole hydrate ("HOBT") (4.0 g, 26.1 mmole) in 40 mL of DMF and 1.6 mL of *N*-Methylmorpholine ("NMM") is charged with 1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide ("EDAC") (2.0 g, 10.44 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and ethyl acetate ("EtOAc"). The organic layer is then washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered, evaporated and chromatographed over flash silica with EtOAc to give 25.

one or more matrix metalloproteases in vivo. The ability of the inhibitors to selectively bind matrix metalloprotease is thus taken advantage of to map the distribution of these enzymes in situ. The techniques can also be employed in histological procedures and the labeled invention compounds can be used in competitive immunoassays.

The following non-limiting examples illustrate the compounds, compositions, and uses of the present invention.

Example 1

Synthesis of (2R)-Isobutyl-3-(N-hydroxycarboxamido)-propanoic acid amide of 1N-(carbomethoxymethyl)-caprolactam-(3S)-amine (1)



(2R)-Isobutyl-3-carbo-*tert*-butoxypropionic acid amide of caprolactam-(3S)-amine (25). A mixture of acid 23 (2.0 g, 8.70 mmole), caprolactam-[3S]-amine 24 (1.23 g, 9.57 mmole) and 1-Hydroxybenzotriazole hydrate ("HOBT") (4.0 g, 26.1 mmole) in 40 mL of DMF and 1.6 mL of *N*-Methylmorpholine ("NMM") is charged with 1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide ("EDAC") (2.0 g, 10.44 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and ethyl acetate ("EtOAc"). The organic layer is then washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered, evaporated and chromatographed over flash silica with EtOAc to give 25.

(2R)-Isobutyl-3-carbo-*tert*-butoxypropanoic acid amide of 1*N*-(carbomethoxy-methyl)-caprolactam-(3*S*)-amine (26). The caprolactam **25** (914 mg, 2.69 mmole) is taken in 10 mL THF and cooled to -78°C under argon. To this is added 1 M lithium bis(trimethylsilyl)amide (2.69 mL, 2.69 mmole) and the reaction stirred for 5 min. Methyl bromoacetate (256 µL, 2.69 mmole) is added and stirred for 2 hr. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1*N* HCl, NaHCO₃, and brine, dried over MgSO₄, filtered and evaporated to give crude material which is then chromatographed over flash silica with hexanes:EtOAc (1:4) to give **26**.

(2R)-Isobutyl-3-carboxypropanoic acid amide of *N*-(carbomethoxymethyl)-caprolactam-(3*S*)-amine (27). Trifluoroacetic acid (3 mL) is added via syringe to a solution of *tert*-butyl ester **26** (380 mg, 0.922 mmole) in 3 mL CH₂Cl₂ under argon and the resulting mixture is stirred for 2 hr at room temperature. The material is then concentrated under vacuum to give **27** which is carried forward without purification.

(2R)-Isobutyl-3-(*O*-benzyl-*N*-hydroxycarboxamido)-propanoic acid amide of 1*N*-(carbomethoxymethyl)-caprolactam-(3*S*)-amine (28). A mixture of acid **27** (345 mg, 0.863 mmole), *O*-benzyl hydroxylamine hydrochloride (166 mg, 1.035 mmole) and HOBT (397 mg, 2.59 mmole) in 5 mL of *N,N*-Dimethylformamide ("DMF") and 260 µL of NMM is charged with EDAC (199 mg, 1.035 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1*N* HCl, NaHCO₃, and brine, dried over MgSO₄, filtered and evaporated to give crude material which is then chromatographed over flash silica with EtOAc to give the title compound **28**.

(2R)-Isobutyl-3-(*N*-hydroxycarboxamido)-propanoic acid amide of 1*N*-(carbomethoxy-methyl)-caprolactam-(3*S*)-amine (1). The benzyl hydroxamic acid **28** (325 mg, 0.705 mmole) is taken in 6 mL EtOH and the mixture is charged with 10% palladium on carbon (60 mg) and stirred under one atmosphere of hydrogen for 45 min. The mixture is then filtered through celite and concentrated to give 240 mg of crude material which is then chromatographed over flash silica with EtOAc:formic acid (98:2) and then recrystallized from hexanes:EtOAc (2:1) to give the pure desired hydroxamic acid **1**.

(2R)-Isobutyl-3-carbo-*tert*-butoxypropanoic acid amide of 1*N*-(carbomethoxymethyl)-caprolactam-(3*S*)-amine (26). The caprolactam **25** (914 mg, 2.69 mmole) is taken in 10 mL THF and cooled to -78°C under argon. To this is added 1 M lithium bis(trimethylsilyl)amide (2.69 mL, 2.69 mmole) and the reaction stirred for 5 min. Methyl bromoacetate (256 µL, 2.69 mmole) is added and stirred for 2 hr. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered and evaporated to give crude material which is then chromatographed over flash silica with hexanes:EtOAc (1:4) to give **26**.


(2R)-Isobutyl-3-carboxypropanoic acid amide of *N*-(carbomethoxymethyl)-caprolactam-(3*S*)-amine (27). Trifluoroacetic acid (3 mL) is added via syringe to a solution of *tert*-butyl ester **26** (380 mg, 0.922 mmole) in 3 mL CH₂Cl₂ under argon and the resulting mixture is stirred for 2 hr at room temperature. The material is then concentrated under vacuum to give **27** which is carried forward without purification.

(2R)-Isobutyl-3-(*O*-benzyl-*N*-hydroxycarboxamido)-propanoic acid amide of 1*N*-(carbomethoxymethyl)-caprolactam-(3*S*)-amine (28). A mixture of acid **27** (345 mg, 0.863 mmole), *O*-benzyl hydroxylamine hydrochloride (166 mg, 1.035 mmole) and HOBT (397 mg, 2.59 mmole) in 5 mL of *N,N*-Dimethylformamide ("DMF") and 260 µL of NMM is charged with EDAC (199 mg, 1.035 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered and evaporated to give crude material which is then chromatographed over flash silica with EtOAc to give the title compound **28**.

(2R)-Isobutyl-3-(*N*-hydroxycarboxamido)-propanoic acid amide of 1*N*-(carbomethoxy-methyl)-caprolactam-(3*S*)-amine (1). The benzyl hydroxamic acid **28** (325 mg, 0.705 mmole) is taken in 6 mL EtOH and the mixture is charged with 10% palladium on carbon (60 mg) and stirred under one atmosphere of hydrogen for 45 min. The mixture is then filtered through celite and concentrated to give 240 mg of crude material which is then chromatographed over flash silica with EtOAc:formic acid (98:2) and then recrystallized from hexanes:EtOAc (2:1) to give the pure desired hydroxamic acid **1**.

Example 2


Synthesis of (2R)-Isobutyl-3-(N-hydroxycarboxamido)-propionic acid amide of 1N-(methyl-carboxamidomethyl)-caprolactam-(3S)-amine (2)



(2R)-Isobutyl-3-(N-hydroxycarboxamido)-propionic acid amide of 1N-(methyl-carboxamidomethyl)-caprolactam-(3S)-amine (2). The methyl ester 1 (80 mg, 0.216 mmole) is taken in 5 mL of 8M methyl amine in MeOH and stirred for 15 hours. The solvent is removed and the residue is chromatographed over flash silica with EtOAc:formic acid (97:3) to give 2.

Example 3

Synthesis of (2R)-Isobutyl-3-(N-hydroxycarboxamido)-propionic acid amide of 1N-(carbo-tert-butoxy-methyl)-caprolactam-(3S)-amine (3)




(2R)-Isobutyl-3-carboxypropionic acid amide of caprolactam-(3S)-amine (29).

Trifluoroacetic acid (15 mL) is added via syringe to a solution of *tert*-butyl ester 25 (2.2 g, 6.47 mmole) in 15 mL CH₂Cl₂ under argon and the resulting mixture is stirred for 2 hr at room temperature. The material is then concentrated under vacuum to give 29 which is carried forward without purification.

(2R)-Isobutyl-3-carbomethoxypropionic acid amide of caprolactam-(3S)-amine (30). To a solution of acid 29 (1.24 g, 4.37 mmole) in 5 mL MeOH is added an excess of diazomethane in ether. The excess diazomethane is then quenched with acetic acid

Example 2


Synthesis of (2R)-Isobutyl-3-(N-hydroxycarboxamido)-propionic acid amide of 1N-(methyl-carboxamidomethyl)-caprolactam-(3S)-amine (2)



(2R)-Isobutyl-3-(N-hydroxycarboxamido)-propionic acid amide of 1N-(methyl-carboxamidomethyl)-caprolactam-(3S)-amine (2). The methyl ester 1 (80 mg, .216 mmole) is taken in 5 mL of 8M methyl amine in MeOH and stirred for 15 hours. The solvent is removed and the residue chromatographed over flash silica with EtOAc:formic acid (97:3) to give 2.

Example 3

Synthesis of (2R)-Isobutyl-3-(N-hydroxycarboxamido)-propionic acid amide of 1N-(carbo-tert-butoxy-methyl)-caprolactam-(3S)-amine (3)



(2R)-Isobutyl-3-carboxypropionic acid amide of caprolactam-(3S)-amine (29).

Trifluoroacetic acid (15 mL) is added via syringe to a solution of *tert*-butyl ester 25 (2.2 g, 6.47 mmole) in 15 mL CH₂Cl₂ under argon and the resulting mixture is stirred for 2 hr at room temperature. The material is then concentrated under vacuum to give 29 which is carried forward without purification.

(2R)-Isobutyl-3-carbomethoxypropionic acid amide of caprolactam-(3S)-amine (30). To a solution of acid 29 (1.24 g, 4.37 mmole) in 5 mL MeOH is added an excess of diazomethane in ether. The excess diazomethane is then quenched with acetic acid


and the solvent is evaporated. The residue is chromatographed over flash silica with EtOAc to give the desired ester 30.

(2R)-Isobutyl-3-carbomethoxypropionic acid amide of 1N-(carbo-*tert*-butoxy-methyl)-caprolactam-(3S)-amine (31). The caprolactam 30 (200 mg, 0.67 mmole) is taken in 5 mL dry tetrahydrofuran ("THF") and cooled to -78°C under argon. To this solution is added 1 M lithium bis(trimethylsilyl)amide (0.67 mL, 0.67 mmole) and the reaction is stirred for 5 min. *tert*-Butyl bromoacetate (99 μL , 0.67 mmole) is added and stirred for 2 hr. The reaction is then partitioned between water and EtOAc. The organic layer is washed with 1N HCl, NaHCO_3 , and brine, dried over MgSO_4 , filtered and evaporated to give a crude material which is chromatographed over flash silica with hexanes:EtOAc (1:4) to give pure desired ester 31.

(2R)-Isobutyl-3-(N-hydroxycarboxamido)-propionic acid amide of 1N-(carbo-*tert*-butoxy-methyl)-caprolactam-(3S)-amine (3). Ester 31 (130 mg, 0.29 mmole) is added to NH_2OK (1.3 mL, 1 eq in MeOH, prepared according to Fieser and Fieser, Reagents for Organic Synthesis, Vol. 1, p. 478 (1967)) and stirred for 24 hr. The solvent is evaporated and the residue is dissolved in 1N HCl and extracted with EtOAc. The organic layer is dried over MgSO_4 , evaporated and the residue is chromatographed over flash silica with EtOAc:formic acid (98:2) to give 3.

Example 4

Synthesis of (2R)-Isobutyl-3-(N-hydroxycarboxamido)-propionic acid amide of 1N-Benzyl-caprolactam-(3S)-amine (4).



(2R)-Isobutyl-3-carbomethoxypropionic acid amide of 1N-Benzylcaprolactam-(3S)-amine (32). The caprolactam 30 (200 mg, 0.67 mmole) is taken in 5 mL dry THF and is cooled to -78°C under argon. To this is added 1 M lithium bis(trimethylsilyl)amide (0.67 mL, 0.67 mmole) and the reaction stirred for 5 min. Benzyl bromide (80 μL , 0.67 mmole) is added and stirred for 2 hr. The reaction is then partitioned between water and EtOAc. The organic layer is washed with 1N HCl, NaHCO_3 , and brine, dried over MgSO_4 , filtered and evaporated to give a crude oil which is chromatographed over flash silica with hexanes:EtOAc (1:4) to give 32.

(2R)-Isobutyl-3-(N-hydroxycarboxamido)-propionic acid amide of 1N-Benzyl-caprolactam-(3S)-amine (4). Ester 32 (200 mg, 0.52 mmole) is added to is added to NH_2OK (1.0 mL, 1 eq in MeOH, prepared according to Fieser and Fieser, Vol 1, p 478)


and the solvent is evaporated. The residue is chromatographed over flash silica with EtOAc to give the desired ester 30.

(2R)-Isobutyl-3-carbomethoxypropionic acid amide of 1N-(carbo-*tert*-butoxy-methyl)-caprolactam-(3S)-amine (31). The caprolactam 30 (200 mg, 0.67 mmole) is taken in 5 mL dry tetrahydrofuran ("THF") and cooled to -78°C under argon. To this solution is added 1 M lithium bis(trimethylsilyl)amide (0.67 mL, 0.67 mmole) and the reaction is stirred for 5 min. *tert*-Butyl bromoacetate (99 μL , 0.67 mmole) is added and stirred for 2 hr. The reaction is then partitioned between water and EtOAc. The organic layer is washed with 1N HCl, NaHCO_3 , and brine, dried over MgSO_4 , filtered and evaporated to give a crude material which is chromatographed over flash silica with hexanes:EtOAc (1:4) to give pure desired ester 31.

(2R)-Isobutyl-3-(*N*-hydroxycarboxamido)-propionic acid amide of 1N-(carbo-*tert*-butoxy-methyl)-caprolactam-(3S)-amine (3). Ester 31 (130 mg, 0.29 mmole) is added to NH_2OK (1.3 mL, 1 eq in MeOH, prepared according to Fieser and Fieser, Reagents for Organic Synthesis, Vol. 1, p. 478 (1967)) and stirred for 24 hr. The solvent is evaporated and the residue is dissolved in 1N HCl and extracted with EtOAc. The organic layer is dried over MgSO_4 , evaporated and the residue is chromatographed over flash silica with EtOAc:formic acid (98:2) to give 3.

Example 4

Synthesis of (2R)-Isobutyl-3-(*N*-hydroxycarboxamido)-propionic acid amide of 1N-Benzyl-caprolactam-(3S)-amine (4).




(2R)-Isobutyl-3-carbomethoxypropionic acid amide of 1N-Benzylcaprolactam-(3S)-amine (32). The caprolactam 30 (200 mg, 0.67 mmole) is taken in 5 mL dry THF and is cooled to -78°C under argon. To this is added 1 M lithium bis(trimethylsilyl)amide (0.67 mL, 0.67 mmole) and the reaction stirred for 5 min. Benzyl bromide (80 μL , 0.67 mmole) is added and stirred for 2 hr. The reaction is then partitioned between water and EtOAc. The organic layer is washed with 1N HCl, NaHCO_3 , and brine, dried over MgSO_4 , filtered and evaporated to give a crude oil which is chromatographed over flash silica with hexanes:EtOAc (1:4) to give 32.

(2R)-Isobutyl-3-(*N*-hydroxycarboxamido)-propionic acid amide of 1N-Benzyl-caprolactam-(3S)-amine (4). Ester 32 (200 mg, 0.52 mmole) is added to is added to NH_2OK (1.0 mL, 1 eq in MeOH, prepared according to Fieser and Fieser, Vol 1, p 478)

and stirred for 24 hr. The solvent is evaporated and the residue is dissolved in 1N HCl and extracted with EtOAc. The organic layer is dried over MgSO₄, evaporated, and the residue is chromatographed over flash silica with EtOAc:formic acid (98:2) to give 4.

Example 5

Synthesis of (2R)-Isobutyl-3-N-hydroxycarboxamidopropionic acid amide of 1N-(carbobenzyloxy-methyl)-caprolactam-(3S)-amine (5)



tert-Butoxycarboxylic acid amide of caprolactam-(3S)-amine (33). A solution of 24 (49.0 g, 383 mmole) in 350 mL DMSO is charged with di-*tert*-butyl dicarbonate (83.5 g, 383 mmole) and stirred for 5 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1N HCl, and brine, dried over MgSO₄, filtered and evaporated. The residue is recrystallized from ether:hexanes (2:1) to give 33.


tert-Butoxycarboxylic acid amide of 1N-(carbobenzyloxymethyl)-caprolactam-(3S)-amine (34). Lithium bis(trimethylsilyl)amide (44.7 mL, 44.7, 1 M in THF) is added to a solution of the caprolactam 33 (10.2 g, 44.7 mmol) in THF (100 mL) at -78° C under argon and stirred for 15 min. Benzyl bromoacetate (7.08 mL, 44.7 mmol) is added to the solution via syringe, warmed to room temperature, and stirred for 2 hr. The reaction is partitioned between H₂O and EtOAc. The organic layer is washed with aqueous NaHCO₃, aqueous NaCl, and dried over MgSO₄. The crude product is chromatographed on flash silica with hexane:EtOAc (1:1) to give 34.

1N-(carbobenzyloxymethyl)-caprolactam-(3S)-amine trifluoroacetic acid salt (35). Trifluoroacetic acid (15 mL) is added via syringe to a solution of *tert*-butyl

and stirred for 24 hr. The solvent is evaporated and the residue is dissolved in 1N HCl and extracted with EtOAc. The organic layer is dried over MgSO₄, evaporated, and the residue is chromatographed over flash silica with EtOAc:formic acid (98:2) to give 4.

Example 5

Synthesis of (2R)-Isobutyl-3-N-hydroxycarboxamidopropionic acid amide of 1N-(carbobenzyloxy-methyl)-caprolactam-(3S)-amine (5)



tert-Butoxycarboxylic acid amide of caprolactam-(3S)-amine (33). A solution of 24 (49.0 g, 383 mmole) in 350 mL DMSO is charged with di-*tert*-butyl dicarbonate (83.5 g, 383 mmole) and stirred for 5 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1N HCl, and brine, dried over MgSO₄, filtered and evaporated. The residue is recrystallized from ether:hexanes (2:1) to give 33.

tert-Butoxycarboxylic acid amide of 1N-(carbobenzyloxymethyl)-caprolactam-(3S)-amine (34). Lithium bis(trimethylsilyl)amide (44.7 mL, 44.7, 1 M in THF) is added to a solution of the caprolactam 33 (10.2 g, 44.7 mmol) in THF (100 mL) at -78° C under argon and stirred for 15 min. Benzyl bromoacetate (7.08 mL, 44.7 mmol) is added to the solution via syringe, warmed to room temperature, and stirred for 2 hr. The reaction is partitioned between H₂O and EtOAc. The organic layer is washed with aqueous NaHCO₃, aqueous NaCl, and dried over MgSO₄. The crude product is chromatographed on flash silica with hexane:EtOAc (1:1) to give 34.

1N-(carbobenzyloxymethyl)-caprolactam-(3S)-amine trifluoroacetic acid salt (35). Trifluoroacetic acid (15 mL) is added via syringe to a solution of *tert*-butyl

carboxamate **34** (5.0 g, 13.2 mmole) in 15 mL under argon and the resulting mixture is stirred for 1 hr at room temperature. The material is then concentrated under vacuum to give **35** which is carried forward without purification.

(2R)-Isobutyl-3-carbo-*tert*-butoxypropionic acid amide of 1N-(carbobenzyloxy-methyl)-caprolactam-(3S)-amine (**36**). A mixture of acid **23** (1.67, 7.25 mmole), caprolactam **35** (2.0 g, 7.25 mmole) and HOBt (2.94 g, 21.75 mmole) in 15 mL of DMF and 1.5 mL of NMM is charged with EDAC (1.67 g, 8.70 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc and then washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered, evaporated and chromatographed over flash silica with EtOAc:hexanes (2:1) to give pure desired amide **36**.


(2R)-Isobutyl-3-carbomethoxypropionic acid amide of 1N-(carbobenzyloxy-methyl)-caprolactam-(3S)-amine (**38**). Trifluoroacetic acid (5 mL) is added via syringe to a solution of *tert*-butyl ester **36** (1.2 g, 2.46 mmole) in 5 mL of CH₂Cl₂ under argon and the resulting mixture is stirred for 2 hr at room temperature. The material is then concentrated under vacuum to give acid **37** as a clear oil which is carried forward without purification.

To a solution of acid **37** (1.06 g, 2.46 mmole) in 5 mL MeOH is added an excess of diazomethane in ether. The excess diazomethane is then quenched with acetic acid and the solvent evaporated. The residue is chromatographed over flash silica with EtOAc:hexanes (2:1) to give the desired ester **38**.

(2R)-Isobutyl-3-N-hydroxycarboxamidopropionic acid amide of 1N-(carbobenzyloxy-methyl)-caprolactam-(3S)-amine (**5**). Ester **38** (420 mg, 0.94 mmole) is added to a solution of NH₂OK (1.3 mL, 1 eq in MeOH, prepared according to Fieser and Fieser, Vol 1, p 478) and stirred for 24 hours. The solvent is then evaporated and the residue dissolved in 1N HCl and extracted with EtOAc. The organic layer is dried over MgSO₄, evaporated and the residue chromatographed over flash silica with EtOAc:formic acid (98:2) to give **5**.

Example 6

Synthesis of (2R)-Isobutyl-3-(N-hydroxycarboxamido)-propionic acid amide of 1N-(N-Benzyl-carboxamidomethyl)-caprolactam-(3S)-amine (**6**)



carboxamate **34** (5.0 g, 13.2 mmole) in 15 mL under argon and the resulting mixture is stirred for 1 hr at room temperature. The material is then concentrated under vacuum to give **35** which is carried forward without purification.

(2R)-Isobutyl-3-carbo-*tert*-butoxypropionic acid amide of 1N-(carbobenzyloxy-methyl)-caprolactam-(3S)-amine (**36**). A mixture of acid **23** (1.67, 7.25 mmole), caprolactam **35** (2.0 g, 7.25 mmole) and HOBT (2.94 g, 21.75 mmole) in 15 mL of DMF and 1.5 mL of NMM is charged with EDAC (1.67 g, 8.70 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc and then washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered, evaporated and chromatographed over flash silica with EtOAc:hexanes (2:1) to give pure desired amide **36**.


(2R)-Isobutyl-3-carbomethoxypropionic acid amide of 1N-(carbobenzyloxy-methyl)-caprolactam-(3S)-amine (**38**). Trifluoroacetic acid (5 mL) is added via syringe to a solution of *tert*-butyl ester **36** (1.2 g, 2.46 mmole) in 5 mL of CH₂Cl₂ under argon and the resulting mixture is stirred for 2 hr at room temperature. The material is then concentrated under vacuum to give acid **37** as a clear oil which is carried forward without purification.


To a solution of acid **37** (1.06 g, 2.46 mmole) in 5 mL MeOH is added an excess of diazomethane in ether. The excess diazomethane is then quenched with acetic acid and the solvent evaporated. The residue is chromatographed over flash silica with EtOAc:hexanes (2:1) to give the desired ester **38**.

(2R)-Isobutyl-3-*N*-hydroxycarboxamidopropionic acid amide of 1N-(carbobenzyloxy-methyl)-caprolactam-(3S)-amine (**5**). Ester **38** (420 mg, 0.94 mmole) is added to a solution of NH₂OK (1.3 mL, 1 eq in MeOH, prepared according to Fieser and Fieser, Vol 1, p 478) and stirred for 24 hours. The solvent is then evaporated and the residue dissolved in 1N HCl and extracted with EtOAc. The organic layer is dried over MgSO₄, evaporated and the residue chromatographed over flash silica with EtOAc:formic acid (98:2) to give **5**.

Example 6

Synthesis of (2R)-Isobutyl-3-(*N*-hydroxycarboxamido)-propionic acid amide of 1N-(*N*-Benzyl-carboxamidomethyl)-caprolactam-(3S)-amine (**6**)






tert-Butoxycarboxylic acid amide of 1*N*-(carbomethoxymethyl)-caprolactam-(3*S*)-amine (39). Lithium bis(trimethylsilyl)amide (83.3 mL, 83.3, 1 M in THF) is added to a solution of the caprolactam 33 (19.0 g, 83.3 mmol) in THF (200 mL) at -78°C under argon and stirred for 15 min. Methyl bromoacetate (7.88 mL, 83.3 mmol) is added to the solution via syringe, warmed to room temperature, and stirred for 1 hr. The reaction is partitioned between H_2O and EtOAc. The organic layer is washed with aqueous NaHCO_3 , aqueous NaCl, and dried over MgSO_4 . The crude product is chromatographed on flash silica with EtOAc to give 39.

tert-Butoxycarboxylic acid amide of 1*N*-(*N*-benzylcarboxamidomethyl)-caprolactam-(3*S*)-amine (40). The methyl ester 39 (2.5 g, 8.33 mmole) is taken in 10 mL MeOH and the mixture is charged with benzyl amine (8.7 mL, 79.7 mmole) and is stirred for 15 hours. The solvent is removed and the residue is chromatographed over flash silica with EtOAc:hexane (1:1) to give 40.

1*N*-(*N*-Benzylcarboxamidomethyl)-caprolactam-(3*S*)-amine hydrochloride (41). The amide 40 (2.2 g, 5.87 mmole) is taken in 50 mL ether at 0°C and dry HCl is bubbled through for 10 min. The solid is filtered and washed with ether to give 41.

(2*R*)-Isobutyl-3-carbo-*tert*-butoxypropionic acid amide of 1*N*-(*N*-benzylcarboxamido-methyl)-caprolactam-(3*S*)-amine (42). A mixture of acid 23 (1.0 g, 4.35 mmole), caprolactam 41 (1.35 g, 4.35 mmole) and HOBt (2.0 g, 13.0 mmole) in 15 mL of DMF and 1.3 mL of NMM is charged with EDAC (1.0 g, 5.22 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1*N* HCl, NaHCO_3 , and brine, dried over MgSO_4 , filtered and evaporated to give 1.9 g of crude solid which is chromatographed over flash silica with EtOAc to give 42.

(2*R*)-Isobutyl-3-carboxypropionic acid amide of 1*N*-(*N*-benzylcarboxamidomethyl)-caprolactam-(3*S*)-amine (43). Trifluoroacetic acid (5 mL) is added via syringe to a solution of *tert*-butyl ester 42 (1.51 g, 3.10 mmole) in 5 mL CH_2Cl_2 under argon and the resulting mixture is stirred for 2 hr at room temperature. The material is



tert-Butoxycarboxylic acid amide of 1*N*-(carbomethoxymethyl)-caprolactam-(3*S*)-amine (39). Lithium bis(trimethylsilyl)amide (83.3 mL, 83.3, 1 M in THF) is added to a solution of the caprolactam 33 (19.0 g, 83.3 mmol) in THF (200 mL) at -78° C under argon and stirred for 15 min. Methyl bromoacetate (7.88 mL, 83.3 mmol) is added to the solution via syringe, warmed to room temperature, and stirred for 1 hr. The reaction is partitioned between H₂O and EtOAc. The organic layer is washed with aqueous NaHCO₃, aqueous NaCl, and dried over MgSO₄. The crude product is chromatographed on flash silica with EtOAc to give 39.

tert-Butoxycarboxylic acid amide of 1*N*-(*N*-benzylcarboxamidomethyl)-caprolactam-(3*S*)-amine (40). The methyl ester 39 (2.5 g, 8.33 mmole) is taken in 10 mL MeOH and the mixture is charged with benzyl amine (8.7 mL, 79.7 mmole) and is stirred for 15 hours. The solvent is removed and the residue is chromatographed over flash silica with EtOAc:hexane (1:1) to give 40.

1*N*-(*N*-Benzylcarboxamidomethyl)-caprolactam-(3*S*)-amine hydrochloride (41). The amide 40 (2.2 g, 5.87 mmole) is taken in 50 mL ether at 0° C and dry HCl is bubbled through for 10 min. The solid is filtered and washed with ether to give 41.

(2*R*)-Isobutyl-3-carbo-*tert*-butoxypropionic acid amide of 1*N*-(*N*-benzylcarboxamido-methyl)-caprolactam-(3*S*)-amine (42). A mixture of acid 23 (1.0 g, 4.35 mmole), caprolactam 41 (1.35 g, 4.35 mmole) and HOBt (2.0 g, 13.0 mmole) in 15 mL of DMF and 1.3 mL of NMM is charged with EDAC (1.0 g, 5.22 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered and evaporated to give 1.9 g of crude solid which is chromatographed over flash silica with EtOAc to give 42.

(2*R*)-Isobutyl-3-carboxypropionic acid amide of 1*N*-(*N*-benzylcarboxamidomethyl)-caprolactam-(3*S*)-amine (43). Trifluoroacetic acid (5 mL) is added via syringe to a solution of *tert*-butyl ester 42 (1.51 g, 3.10 mmole) in 5 mL CH₂Cl₂ under argon and the resulting mixture is stirred for 2 hr at room temperature. The material is

then concentrated under vacuum to give **43** which is carried forward without purification.


(2R)-Isobutyl-3-carbomethoxypropionic acid amide of 1N-(N-benzylcarbox-amido-methyl)-caprolactam-(3S)-amine (44). To a solution of acid **43** (1.51 g, 3.50 mmole) in 5 mL MeOH is added an excess of diazomethane in ether. The excess diazomethane is quenched with acetic acid and the solvent evaporated. The residue is chromatographed over flash silica with EtOAc:hexanes (2:1) to give **44**.

(2R)-Isobutyl-3-(N-hydroxycarboxamido)-propionic acid amide of 1N-(N-benzylcarboxamidomethyl)-caprolactam-(3S)-amine (6). Ester **44** (800 mg, 1.80 mmole) is added to a solution of NH₂OK (1.5 mL, 1 eq in MeOH, prepared according to Fieser and Fieser, Vol 1, p 478) and stirred for 24 hours. The solvent is evaporated and the residue is dissolved in 1N HCl and extracted with EtOAc. The organic layer is dried over MgSO₄, filtered, evaporated and the residue is chromatographed over flash silica with EtOAc:formic acid (97:3) to give **6**.

then concentrated under vacuum to give 43 which is carried forward without purification.

(2R)-Isobutyl-3-carbomethoxypropionic acid amide of 1N-(N-benzylcarbox-amido-methyl)-caprolactam-(3S)-amine (44). To a solution of acid 43 (1.51 g, 3.50 mmole) in 5 mL MeOH is added an excess of diazomethane in ether. The excess diazomethane is quenched with acetic acid and the solvent evaporated. The residue is chromatographed over flash silica with EtOAc:hexanes (2:1) to give 44.

(2R)-Isobutyl-3-(N-hydroxycarboxamido)-propionic acid amide of 1N-(N-benzylcarboxamidomethyl)-caprolactam-(3S)-amine (6). Ester 44 (800 mg, 1.80 mmole) is added to a solution of NH₂OK (1.5 mL, 1 eq in MeOH, prepared according to Fieser and Fieser, Vol 1, p 478) and stirred for 24 hours. The solvent is evaporated and the residue is dissolved in 1N HCl and extracted with EtOAc. The organic layer is dried over MgSO₄, filtered, evaporated and the residue is chromatographed over flash silica with EtOAc:formic acid (97:3) to give 6.


Example 7Synthesis of (2R)-Isobutyl-3-N-hydroxycarboxamidopropionic acid amide of 1N-(n-butylcarboxamidomethyl)-caprolactam-(3S)-amine (7)

tert-Butoxycarboxylic acid amide of 1N-(n-butylcarboxamidomethyl)-caprolactam-(3S)-amine (45). The methyl ester 39 (2.5 g, 8.33 mmole) is taken in 10 mL MeOH and the mixture is charged with butyl amine (8.1 mL, 79.7 mmole) and stirred for 15 hr. The solvent is removed and the residue is chromatographed over flash silica with EtOAc:hexane (2:1) to give 45.

1N-(n-butylcarboxamidomethyl)-(3S)-aminocaprolactam hydrochloride (46). The amide 45 (2.2 g, 6.43 mmole) is taken in 50 mL ether at 0° C and dry HCl is bubbled through for 10 min. The solid is filtered and washed with ether to give 46.

(2R)-Isobutyl-(3)-carbo-tert-butoxypropionic acid amide of (1N)-n-butylcarboxamido-methyl-caprolactam-(3S)-amine (47). A mixture of acid 23, (1.76 g, 7.65 mmole), 1N-(n-butylcarboxamidomethyl)-caprolactam-(3S)-amine 46 (3.17 g, 11.4 mmole) and HOBt (3.08g, 22.80 mmole) in 20 mL of DMF and 2.36 mL of NMM is charged with EDAC (1.74 g, 9.07 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered and evaporated to give 4.68 g of crude solid which is chromatographed over flash silica with EtOAc to give 47.

(2R)-Isobutyl-(3)-carboxypropionic acid amide of (1N)-n-butylcarboxamidomethyl-caprolactam-(3S)-amine (48). Trifluoroacetic acid (15 mL) is added via syringe to a solution of the tert-butyl ester 47 (2.45 g, 5.40 mmole) in 15 mL of CH₂Cl₂ under argon and the resulting mixture is stirred for 1 hr at room temperature. The material is then concentrated under vacuum to give 48 which is carried forward without purification.

Example 7Synthesis of (2R)-Isobutyl-3-N-hydroxycarboxamidopropionic acid amide of 1N-(n-butylcarboxamidomethyl)-caprolactam-(3S)-amine (7)

tert-Butoxycarboxylic acid amide of 1N-(n-butylcarboxamidomethyl)-caprolactam-(3S)-amine (45). The methyl ester 39 (2.5 g, 8.33 mmole) is taken in 10 mL MeOH and the mixture is charged with butyl amine (8.1 mL, 79.7 mmole) and stirred for 15 hr. The solvent is removed and the residue is chromatographed over flash silica with EtOAc:hexane (2:1) to give 45.

1N-(n-butylcarboxamidomethyl)-(3S)-aminocaprolactam hydrochloride (46). The amide 45 (2.2 g, 6.43 mmole) is taken in 50 mL ether at 0° C and dry HCl is bubbled through for 10 min. The solid is filtered and washed with ether to give 46.

(2R)-Isobutyl-(3)-carbo-tert-butoxypropanoic acid amide of (1N)-n-butylcarboxamido-methyl-caprolactam-(3S)-amine (47). A mixture of acid 23, (1.76 g, 7.65 mmole), 1N-(n-butylcarboxamidomethyl)-caprolactam-(3S)-amine 46 (3.17 g, 11.4 mmole) and HOBt (3.08g, 22.80 mmole) in 20 mL of DMF and 2.36 mL of NMM is charged with EDAC (1.74 g, 9.07 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered and evaporated to give 4.68 g of crude solid which is chromatographed over flash silica with EtOAc to give 47.

(2R)-Isobutyl-(3)-carboxypropanoic acid amide of (1N)-n-butylcarboxamidomethyl-caprolactam-(3S)-amine (48). Trifluoroacetic acid (15 mL) is added via syringe to a solution of the tert-butyl ester 47 (2.45 g, 5.40 mmole) in 15 mL of CH₂Cl₂ under argon and the resulting mixture is stirred for 1 hr at room temperature. The material is then concentrated under vacuum to give 48 which is carried forward without purification.

(2R)-Isobutyl-3-carbomethoxypropionic acid amide of 1*N*-(*n*-butylcarboxamidomethyl)-caprolactam-(3*S*)-amine (49). The acid 48 (1.23 g, 3.09 mmole) is taken in methanol and treated with an excess of diazomethane in ether at 0°C, and stirred for one hour. Formic acid (3 drops) is added and the mixture is evaporated to dryness to give 49 which is carried forward without purification.


(2R)-Isobutyl-3-*N*-hydroxycarboxamidopropionic acid amide of 1*N*-(*n*-butylcarboxamidomethyl)-caprolactam-(3*S*)-amine (7). The methyl ester 49 (500 mg, 1.21 mmole) is added to a solution of NH₂OK (1.4 mL, 1 eq in MeOH, prepared according to Fieser and Fieser, Vol 1, p 478) and stirred for 15 hr. The reaction mixture is acidified with acetic acid to pH = 2. The reaction is then partitioned between ethyl acetate and water. The organic layer is then washed with brine, and dried over magnesium sulfate, filtered and evaporated to give a crude solid which is chromatographed with ethyl acetate:water:acetic acid (16:1:1) and then recrystallized from ethyl acetate to give 7.

(2R)-Isobutyl-3-carbomethoxypropionic acid amide of 1*N*-(*n*-butylcarboxamidomethyl)-caprolactam-(3*S*)-amine (49). The acid 48 (1.23 g, 3.09 mmole) is taken in methanol and treated with an excess of diazomethane in ether at 0°C, and stirred for one hour. Formic acid (3 drops) is added and the mixture is evaporated to dryness to give 49 which is carried forward without purification.

(2R)-Isobutyl-3-*N*-hydroxycarboxamidopropionic acid amide of 1*N*-(*n*-butylcarboxamidomethyl)-caprolactam-(3*S*)-amine (7). The methyl ester 49 (500 mg, 1.21 mmole) is added to a solution of NH₂OK (1.4 mL, 1 eq in MeOH, prepared according to Fieser and Fieser, Vol 1, p 478) and stirred for 15 hr. The reaction mixture is acidified with acetic acid to pH = 2. The reaction is then partitioned between ethyl acetate and water. The organic layer is then washed with brine, and dried over magnesium sulfate, filtered and evaporated to give a crude solid which is chromatographed with ethyl acetate:water:acetic acid (16:1:1) and then recrystallized from ethyl acetate to give 7.

Example 8


Synthesis of (2R)-N-Hydroxycarboxamidomethyl heptanoic acid amide of N-(carbomethoxymethyl)-caprolactam-(3S)-amine (8)



3-(1-Oxoheptyl)-(4S)-phenylmethyl-2-oxazolidinone (52). *n*-Butyl lithium (58 mL, 2.5 M in hexanes, 142 mmol) is added to a solution of (S)-4-benzyl-2-oxazolidinone 51 (25 g, 141 mmol) in THF (250 mL) at -78°C under argon and stirred for 15 min. Heptanoyl chloride 50 (21 g, 141 mmol) is added to the solution dropwise

Example 8

Synthesis of (2R)-N-Hydroxycarboxamidomethyl heptanoic acid amide of N-(carbomethoxymethyl)-caprolactam-(3S)-amine (8)



3-(1-Oxoheptyl)-(4S)-phenylmethyl-2-oxazolidinone (52). *n*-Butyl lithium (58 mL, 2.5 M in hexanes, 142 mmol) is added to a solution of (S)-4-benzyl-2-oxazolidinone 51 (25 g, 141 mmol) in THF (250 mL) at -78°C under argon and stirred for 15 min. Heptanoyl chloride 50 (21 g, 141 mmol) is added to the solution dropwise

and stirred for 40 min, then warmed to 0° C for 2 hours. The reaction is quenched with NH₄Cl and extracted with EtOAc. The organic layer is washed with 1 N HCl, aqueous NaHCO₃, aqueous NaCl, and dried over MgSO₄. The crude product is recrystallized from hexane to give 52.

3-[1-Oxo-(2R)-(carbo-*tert*-butoxymethyl)-heptyl]-(4S)-phenylmethyl-2-oxazolidinone (53). Lithium bis(trimethylsilyl)amide (132 mL, 132 mmol, 1 M in THF) is added to a solution of the oxazolidinone 52 (38.0 g, 132 mmol) in THF (100 mL) at -78° C under argon and stirred for 15 min. *tert*-Butyl bromoacetate (26 mL, 132 mmol) is added to the solution via syringe and stirred for 3 hr. The reaction is then warmed to 0° C and stirred for 1.5 hr. The reaction is quenched with NH₄Cl and extracted with EtOAc. The organic layer is washed with 1 N HCl, aqueous NaHCO₃, aqueous NaCl, and dried over MgSO₄. The crude product is recrystallized from hexane:EtOAc (2:1) to give 53.

tert-Butyl-[(3R)-carboxy]octanoate (54). The oxazolidinone 53 (10.0 g, 24.9 mmole) is dissolved in THF/H₂O (100 mL : 25 mL) under argon and cooled to 0° C. Hydrogen peroxide (12 mL, 30%, 106.7 mmol) is added dropwise to the solution, followed by lithium hydroxide monohydrate (1.8 g, 43.9 mmol) in H₂O (40 mL). The reaction is stirred for 3 hr, at which time sodium sulfite (10 g in 40 mL H₂O) is added dropwise and stirred for 20 min. The solution is extracted 3 times with CH₂Cl₂. The organic extracts are combined, washed with aqueous NaCl, dried over MgSO₄, filtered, and the solvent removed to give 54.

1*N*-(carbomethoxymethyl)-caprolactam-(3S)-amine trifluoroacetic acid salt (55). Trifluoroacetic acid (15 mL) is added via syringe to a solution of *tert*-butyl carboxamate 39 (10.0 g, 33.3 mmole) in 15 mL under argon and the resulting mixture is stirred for 1 hr at room temperature. The material is then concentrated under vacuum to give 55 which is carried forward without purification.

(2R)-Carbo-*tert*-butoxymethyl heptanoic acid amide of *N*-carbomethoxymethyl-caprolactam-(3S)-amine (56). The amine 55 (2.16 g, 10.8 mmole) is mixed with the acid 54 (2.64 g, 10.8 mmole), HOBT (4.4 g, 32.4 mmole), EDAC (2.69 g, 14 mmole) in 15 mL of DMF and 2.4 mL (21.6 mmole) of NMM at 0° C, and stirred for 17 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is washed with 1N HCl, H₂O, 1N NaOH, and brine, dried over MgSO₄, filtered and evaporated to give an oil (3.2 g) which is chromatographed over flash silica with hexane:EtOAc (1:1) to give 56.

(2R)-*O*-Benzyl-*N*-hydroxycarboxamidomethyl heptanoic acid amide of *N*-carbomethoxymethyl-caprolactam-(3S)-amine (58). The ester 56 (1.2 g, 2.64 mmole) is dissolved in 15 mL of methylene chloride and cooled to 0° C and trifluoroacetic acid

and stirred for 40 min, then warmed to 0° C for 2 hours. The reaction is quenched with NH₄Cl and extracted with EtOAc. The organic layer is washed with 1 N HCl, aqueous NaHCO₃, aqueous NaCl, and dried over MgSO₄. The crude product is recrystallized from hexane to give 52.

3-[1-Oxo-(2R)-(carbo-*tert*-butoxymethyl)-heptyl]-(4S)-phenylmethyl-2-oxazolidinone (53). Lithium bis(trimethylsilyl)amide (132 mL, 132 mmol, 1 M in THF) is added to a solution of the oxazolidinone 52 (38.0 g, 132 mmol) in THF (100 mL) at -78° C under argon and stirred for 15 min. *tert*-Butyl bromoacetate (26 mL, 132 mmol) is added to the solution via syringe and stirred for 3 hr. The reaction is then warmed to 0° C and stirred for 1.5 hr. The reaction is quenched with NH₄Cl and extracted with EtOAc. The organic layer is washed with 1 N HCl, aqueous NaHCO₃, aqueous NaCl, and dried over MgSO₄. The crude product is recrystallized from hexane:EtOAc (2:1) to give 53.

tert-Butyl-[(3R)-carboxy]octanoate (54). The oxazolidinone 53 (10.0 g, 24.9 mmole) is dissolved in THF/H₂O (100 mL : 25 mL) under argon and cooled to 0° C. Hydrogen peroxide (12 mL, 30%, 106.7 mmol) is added dropwise to the solution, followed by lithium hydroxide monohydrate (1.8 g, 43.9 mmol) in H₂O (40 mL). The reaction is stirred for 3 hr, at which time sodium sulfite (10 g in 40 mL H₂O) is added dropwise and stirred for 20 min. The solution is extracted 3 times with CH₂Cl₂. The organic extracts are combined, washed with aqueous NaCl, dried over MgSO₄, filtered, and the solvent removed to give 54.

1*N*-(carbomethoxymethyl)-caprolactam-(3S)-amine trifluoroacetic acid salt (55). Trifluoroacetic acid (15 mL) is added via syringe to a solution of *tert*-butyl carboxamate 39 (10.0 g, 33.3 mmole) in 15 mL under argon and the resulting mixture is stirred for 1 hr at room temperature. The material is then concentrated under vacuum to give 55 which is carried forward without purification.

(2R)-Carbo-*tert*-butoxymethyl heptanoic acid amide of *N*-carbomethoxymethyl-caprolactam-(3S)-amine (56). The amine 55 (2.16 g, 10.8 mmole) is mixed with the acid 54 (2.64 g, 10.8 mmole), HOBT (4.4 g, 32.4 mmole), EDAC (2.69 g, 14 mmole) in 15 mL of DMF and 2.4 mL (21.6 mmole) of NMM at 0°C, and stirred for 17 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is washed with 1N HCl, H₂O, 1N NaOH, and brine, dried over MgSO₄, filtered and evaporated to give an oil (3.2 g) which is chromatographed over flash silica with hexane:EtOAc (1:1) to give 56.

(2R)-*O*-Benzyl-*N*-hydroxycarboxamidomethyl heptanoic acid amide of *N*-carbomethoxymethyl-caprolactam-(3S)-amine (58). The ester 56 (1.2 g, 2.64 mmole) is dissolved in 15 mL of methylene chloride and cooled to 0°C and trifluoroacetic acid


(15 mL) is added slowly. The mixture is stirred for 1.5 hr at room temperature and evaporated to **57**.

A mixture of the crude acid **57** (1.5 g, 2.64 mmole), *O*-benzyl hydroxylamine hydrochloride (0.51 g, 3.17 mmole) and HOBT (1.07 g, 7.92 mmole) in 10 mL of DMF and 0.99 mL of NMM (9.0 mmole) is charged with EDAC (0.61 g, 3.17 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered and evaporated to give an oil which is chromatographed over flash silica with EtOAc to give **58**.

(2R)-N-Hydroxycarboxamidomethyl heptanoic acid amide of N-(carbomethoxymethyl)-caprolactam-(3S)-amine (8). The benzyl hydroxamic acid **58** (370 mg, 0.78 mmole) is taken in 10 mL of EtOH and the mixture is charged with 10% palladium on carbon (44 mg) and stirred under one atmosphere of hydrogen for 3 hr. The mixture is then filtered through celite and concentrated to an oil. The crude product is crystallized from ethyl acetate to give **8**.

Example 9

Synthesis of (2R)-N-hydroxycarboxamidomethyldecanoic acid amide of 1N-(carbomethoxymethyl)-caprolactam-(3S)-amine (9)



3-(1-Oxodecyl)-(4S)-phenylmethyl-2-oxazolidinone (60). *n*-Butyl lithium (160 mL, 398 mmol, 2.5 M in hexanes) is added to a solution of (S)-4-benzyl-2-oxazolidinone **51** (64 g, 362 mmol) in THF (750 mL) at -78° C under argon and stirred for 15 min. Decanoyl chloride **59** (69 g, 362 mmol) is added to the solution dropwise and stirred for 40 min, then warmed to 0° C for 2 hours. The reaction is quenched with NH₄Cl and extracted with EtOAc. The organic layer is washed with 1 N HCl, aqueous NaHCO₃, aqueous NaCl, and dried over MgSO₄. The crude product is recrystallized from hexane to give **60**.


(15 mL) is added slowly. The mixture is stirred for 1.5 hr at room temperature and evaporated to **57**.

A mixture of the crude acid **57** (1.5 g, 2.64 mmole), *O*-benzyl hydroxylamine hydrochloride (0.51 g, 3.17 mmole) and HOBT (1.07 g, 7.92 mmole) in 10 mL of DMF and 0.99 mL of NMM (9.0 mmole) is charged with EDAC (0.61 g, 3.17 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered and evaporated to give an oil which is chromatographed over flash silica with EtOAc to give **58**.

(2R)-N-Hydroxycarboxamidomethyl heptanoic acid amide of N-(carbomethoxymethyl)-caprolactam-(3S)-amine (8). The benzyl hydroxamic acid **58** (370 mg, 0.78 mmole) is taken in 10 mL of EtOH and the mixture is charged with 10% palladium on carbon (44 mg) and stirred under one atmosphere of hydrogen for 3 hr. The mixture is then filtered through celite and concentrated to an oil. The crude product is crystallized from ethyl acetate to give **8**.

Example 9

Synthesis of (2R)-N-hydroxycarboxamidomethyldecanoic acid amide of 1N-(carbomethoxymethyl)-caprolactam-(3S)-amine (9)



3-(1-Oxodecyl)-(4S)-phenylmethyl-2-oxazolidinone (60). *n*-Butyl lithium (160 mL, 398 mmol, 2.5 M in hexanes) is added to a solution of (S)-4-benzyl-2-oxazolidinone **51** (64 g, 362 mmol) in THF (750 mL) at -78° C under argon and stirred for 15 min. Decanoyl chloride **59** (69 g, 362 mmol) is added to the solution dropwise and stirred for 40 min, then warmed to 0° C for 2 hours. The reaction is quenched with NH₄Cl and extracted with EtOAc. The organic layer is washed with 1 N HCl, aqueous NaHCO₃, aqueous NaCl, and dried over MgSO₄. The crude product is recrystallized from hexane to give **60**.

3-[1-Oxo-(2R)-(carbo-*tert*-butoxymethyl)-decyl]-(4S)-phenylmethyl-2-oxazolidinone (61). Lithium bis(trimethylsilyl)amide (210 mL, 1 M in THF, 210 mmol) is added to a solution of the oxazolidinone **60** (66.6 g, 200 mmol) in THF (100 mL) at -78° C under argon and stirred for 15 min. *tert*-Butyl bromoacetate (31 mL, 200 mmol) is added to the solution via syringe and stirred for 3 hr. The reaction is warmed to 0° C and stirred for 1.5 hr. The reaction is quenched with NH₄Cl and extracted with EtOAc. The organic layer is washed with 1 N HCl, aqueous NaHCO₃, aqueous NaCl, and dried over MgSO₄. The crude product is chromatographed on flash silica with hexane:EtOAc (7:1) to give **61**.

tert-Butyl-[3R-carboxy]undecanoate (62). Oxazolidinone **61** is dissolved in THF/H₂O (100 mL:25 mL) under argon and cooled to 0° C. Hydrogen peroxide (13 mL, 30%, 115.6 mmol) is added dropwise to the solution, followed by lithium hydroxide monohydrate (1.8 g, 43.9 mmol) in H₂O (40 mL). The reaction is stirred for 3 hr, at which time sodium sulfite (10 g in 40 mL H₂O) is added dropwise and stirred 20 min. The solution is extracted 3 times with CH₂Cl₂. The organic extracts are combined, washed with aqueous NaCl, and dried over MgSO₄. The product is purified on a silica gel column using EtOAc as the eluent to give **62**.

(2R)-Carbo-*tert*-butoxymethyldecanoic acid amide of 1*N*-(carbomethoxymethyl)-caprolactam-(3S)-amine (63). A mixture of acid **62** (2.5 g, 8.74 mmole) and 0.9 mL NMM is cooled to -20° C and charged with isobutyl chloroformate (1.45 mL, 8.74 mmole) and stirred for 10 min. The amine **55** (1.75 g, 8.74 mmole) and 0.9 mL NMM in 2 mL DMF is added and stirred for 30 min. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered and evaporated to give 2.8 g of crude solid which chromatographed over flash silica with EtOAc:hexanes (1:1) to give **63**.

(2R)-*O*-Benzyl-*N*-hydroxycarboxamidemethyldecanoic acid amide of 1*N*-(carbomethoxymethyl)-caprolactam-(3S)-amine (65). Trifluoroacetic acid (5 mL) is added via syringe to a solution of *tert*-butyl ester **63** (1.2 g, 2.56 mmole) in 5 mL CH₂Cl₂ under argon and the resulting mixture is stirred for 2 hr at room temperature. The material is then concentrated under vacuum to give **64** which is carried forward without purification.

A mixture of acid **64** (1.05 g, 2.55 mmole), *O*-benzyl hydroxylamine hydrochloride (165 mg, 2.55 mmole) and HOBT (1.2 g, 7.65 mmole) in 15 mL of DMF and 6 mL of NMM is charged with EDAC (572 mg, 2.55 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1N HCl, NaHCO₃, and brine, dried

3-[1-Oxo-(2R)-(carbo-*tert*-butoxymethyl)-decyl]-(4S)-phenylmethyl-2-oxazolidinone (61). Lithium bis(trimethylsilyl)amide (210 mL, 1 M in THF, 210 mmol) is added to a solution of the oxazolidinone **60** (66.6 g, 200 mmol) in THF (100 mL) at -78° C under argon and stirred for 15 min. *tert*-Butyl bromoacetate (31 mL, 200 mmol) is added to the solution via syringe and stirred for 3 hr. The reaction is warmed to 0° C and stirred for 1.5 hr. The reaction is quenched with NH₄Cl and extracted with EtOAc. The organic layer is washed with 1 N HCl, aqueous NaHCO₃, aqueous NaCl, and dried over MgSO₄. The crude product is chromatographed on flash silica with hexane:EtOAc (7:1) to give **61**.

tert-Butyl-[3R-carboxy]undecanoate (62). Oxazolidinone **61** is dissolved in THF/H₂O (100 mL:25 mL) under argon and cooled to 0° C. Hydrogen peroxide (13 mL, 30%, 115.6 mmol) is added dropwise to the solution, followed by lithium hydroxide monohydrate (1.8 g, 43.9 mmol) in H₂O (40 mL). The reaction is stirred for 3 hr, at which time sodium sulfite (10 g in 40 mL H₂O) is added dropwise and stirred 20 min. The solution is extracted 3 times with CH₂Cl₂. The organic extracts are combined, washed with aqueous NaCl, and dried over MgSO₄. The product is purified on a silica gel column using EtOAc as the eluent to give **62**.

(2R)-Carbo-*tert*-butoxymethyldecanoic acid amide of 1*N*-(carbomethoxymethyl)-caprolactam-(3S)-amine (63). A mixture of acid **62** (2.5 g, 8.74 mmole) and 0.9 mL NMM is cooled to -20° C and charged with isobutyl chloroformate (1.45 mL, 8.74 mmole) and stirred for 10 min. The amine **55** (1.75 g, 8.74 mmole) and 0.9 mL NMM in 2 mL DMF is added and stirred for 30 min. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered and evaporated to give 2.8 g of crude solid which chromatographed over flash silica with EtOAc:hexanes (1:1) to give **63**.

(2R)-*O*-Benzyl-*N*-hydroxycarboxamidemethyldecanoic acid amide of 1*N*-(carbomethoxymethyl)-caprolactam-(3S)-amine (65). Trifluoroacetic acid (5 mL) is added via syringe to a solution of *tert*-butyl ester **63** (1.2 g, 2.56 mmole) in 5 mL CH₂Cl₂ under argon and the resulting mixture is stirred for 2 hr at room temperature. The material is then concentrated under vacuum to give **64** which is carried forward without purification.


A mixture of acid **64** (1.05 g, 2.55 mmole), *O*-benzyl hydroxylamine hydrochloride (165 mg, 2.55 mmole) and HOBT (1.2 g, 7.65 mmole) in 15 mL of DMF and 6 mL of NMM is charged with EDAC (572 mg, 2.55 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1N HCl, NaHCO₃, and brine, dried

over MgSO_4 , filtered, evaporated and chromatographed over flash silica with EtOAc to give 65.

(2R)-N-hydroxycarboxamidemethyldecanoic acid amide of 1N-(carbomethoxymethyl)-caprolactam-(3S)-amine (9). The benzyl hydroxamic acid 65 (428 mg, .828 mmole) is taken in 15 mL EtOH and the mixture is charged with 10% palladium on carbon (60 mg) and stirred under one atmosphere of hydrogen for 45 min. The mixture is then filtered through celite and concentrated and chromatographed over flash silica with EtOAc:formic acid (99:1) and then recrystallized from EtOAc to give 9.

Example 10

Synthesis of (2R)-N-Hydroxycarboxamidomethyl decanoic acid amide of N-p-toluenesulfonyl-caprolactam-(3S)-amine (10)



(3S)-tert-Butoxycarbonylamino-N-toluenesulfonyl-caprolactam (66). (3S)-tert-Butoxycarbonylamino caprolactam 33 (2.0 g, 8.8 mmole) is dissolved in THF (20 mL) and cooled to 0°C. A solution of lithium bis(trimethylsilyl)amide (10.6 mL, 10.6 mmole, 1M in THF) is added dropwise. After 15 min, toluenesulfonyl chloride (2.0 g, 10.56 mmole) is added. The resulting mixture is stirred at 0°C for 10 min and room temperature for 30 min. The reaction is quenched by water and extracted by ethyl acetate. The organic layer is then washed with 0.1N HCl, H₂O, and brine, dried over MgSO_4 , filtered and evaporated to give an oil which is chromatographed over flash silica with hexanes:EtOAc (3:1) to give 66.


(2R)-Carbo-tert-butoxymethyl decanoic acid amide of N-toluenesulfonyl-caprolactam-(3S)-amine (68). (3S)-tert-Butoxycarbonylamino-N-toluenesulfonyl-caprolactam 66 (2.2 g, 5.8 mmole) is dissolved in 10 mL of methylene chloride and trifluoroacetic acid (10 mL) is added slowly. The mixture is stirred for 30 min. and evaporated to give 67.

over MgSO₄, filtered, evaporated and chromatographed over flash silica with EtOAc to give **65**.

(2R)-N-hydroxycarboxamidomethyldecanoic acid amide of 1N-(carbomethoxymethyl)-caprolactam-(3S)-amine (9). The benzyl hydroxamic acid **65** (428 mg, .828 mmole) is taken in 15 mL EtOH and the mixture is charged with 10% palladium on carbon (60 mg) and stirred under one atmosphere of hydrogen for 45 min. The mixture is then filtered through celite and concentrated and chromatographed over flash silica with EtOAc:formic acid (99:1) and then recrystallized from EtOAc to give **9**.

Example 10

Synthesis of (2R)-N-Hydroxycarboxamidomethyl decanoic acid amide of N-p-toluenesulfonyl-caprolactam-(3S)-amine (10)



(3S)-tert-Butoxycarbonylamino-N-toluenesulfonyl-caprolactam (66). (3S)-tert-Butoxycarbonyl-amino caprolactam **33** (2.0 g, 8.8 mmole) is dissolved in THF (20 mL) and cooled to 0°C. A solution of lithium bis(trimethylsilyl)amide (10.6 mL, 10.6 mmole, 1M in THF) is added dropwise. After 15 min, toluenesulfonyl chloride (2.0 g, 10.56 mmole) is added. The resulting mixture is stirred at 0°C for 10 min and room temperature for 30 min. The reaction is quenched by water and extracted by ethyl acetate. The organic layer is then washed with 0.1N HCl, H₂O, and brine, dried over MgSO₄, filtered and evaporated to give an oil which is chromatographed over flash silica with hexanes:EtOAc (3:1) to give **66**.

(2R)-Carbo-tert-butoxymethyl decanoic acid amide of N-toluenesulfonyl-caprolactam-(3S)-amine (68). (3S)-tert-Butoxycarbonylamino-N-toluenesulfonyl-caprolactam **66** (2.2 g, 5.8 mmole) is dissolved in 10 mL of methylene chloride and trifluoroacetic acid (10 mL) is added slowly. The mixture is stirred for 30 min. and evaporated to give **67**.

The caprolactam amine **67** (1.6 g, 5.7 mmole) is mixed with **62**, HOBT (2.31 g, 17.1 mmole), EDAC (1.34 g, 7 mmole) in 25 mL of DMF and 1.26 mL (11.4 mmole) of NMM at 0°C, and stirred for 17 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered and evaporated to give an oil (3.1 g) which is chromatographed over flash silica with EtOAc to give **68**.


(2R)-O-Benzyl-N-hydroxycarboxamidomethyl decanoic acid amide of N-toluenesulfonyl-caprolactam-(3S)-amine (70). (3S)-*tert*-Butoxycarbonylamino-N-toluenesulfonyl-caprolactam **68** (2.1 g, 3.8 mmole) is dissolved in 10 mL of methylene chloride, cooled to 0° C, and trifluoroacetic acid (10 mL) is added slowly. The mixture is stirred for 1.5 hr at room temperature and evaporated to give **69**.

A mixture of the crude acid **69** (2 g, 3.8 mmole), *O*-benzyl hydroxylamine hydrochloride (0.73 g, 4.56 mmole) and HOBT (1.54 g, 11.4 mmole) in 20 mL of DMF and 1.4 mL of NMM (12.9 mmole) is charged with EDAC (0.87g, 4.56 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is washed with 1N HCl, 1N NaOH, and brine, dried over MgSO₄, filtered and evaporated to give a semisolid (2.2 g) which is chromatographed over flash silica with EtOAc:hexane (1:1) to give **70**.

(2R)-N-Hydroxycarboxamidomethyl decanoic acid amide of N-toluenesulfonyl-caprolactam-(3S)-amine (10). The benzyl hydroxamic acid **70** (680 mg, 1.13 mmole) is taken in 10 mL of EtOH and the mixture is charged with 10% palladium on carbon (68 mg) and stirred under one atmosphere of hydrogen for 2.5 hr. The mixture is filtered through celite and concentrated to give a solid, which is purified by flash chromatography (silica, EtOAc) to give **10**.

Example 11

Synthesis of (2R)-N-Hydroxycarboxamidomethyl decanoic acid amide of N-[(2-methoxy)ethyl]-caprolactam-(3S)-amine (11)



The caprolactam amine **67** (1.6 g, 5.7 mmole) is mixed with **62**, HOBT (2.31 g, 17.1 mmole), EDAC (1.34 g, 7 mmole) in 25 mL of DMF and 1.26 mL (11.4 mmole) of NMM at 0°C, and stirred for 17 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered and evaporated to give an oil (3.1 g) which is chromatographed over flash silica with EtOAc to give **68**.


(2R)-O-Benzyl-N-hydroxycarboxamidomethyl decanoic acid amide of N-toluenesulfonyl-caprolactam-(3S)-amine (70). (3S)-*tert*-Butoxycarbonylamino-N-toluenesulfonyl-caprolactam **68** (2.1 g, 3.8 mmole) is dissolved in 10 mL of methylene chloride, cooled to 0° C, and trifluoroacetic acid (10 mL) is added slowly. The mixture is stirred for 1.5 hr at room temperature and evaporated to give **69**.


A mixture of the crude acid **69** (2 g, 3.8 mmole), *O*-benzyl hydroxylamine hydrochloride (0.73 g, 4.56 mmole) and HOBT (1.54 g, 11.4 mmole) in 20 mL of DMF and 1.4 mL of NMM (12.9 mmole) is charged with EDAC (0.87g, 4.56 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is washed with 1N HCl, 1N NaOH, and brine, dried over MgSO₄, filtered and evaporated to give a semisolid (2.2 g) which is chromatographed over flash silica with EtOAc:hexane (1:1) to give **70**.

(2R)-N-Hydroxycarboxamidomethyl decanoic acid amide of N-toluenesulfonyl-caprolactam-(3S)-amine (10). The benzyl hydroxamic acid **70** (680 mg, 1.13 mmole) is taken in 10 mL of EtOH and the mixture is charged with 10% palladium on carbon (68 mg) and stirred under one atmosphere of hydrogen for 2.5 hr. The mixture is filtered through celite and concentrated to give a solid, which is purified by flash chromatography (silica, EtOAc) to give **10**.

Example 11

Synthesis of 2R)-N-Hydroxycarboxamidomethyl decanoic acid amide of N-[(2-methoxy)ethyl]-caprolactam-(3S)-amine (11)





(3S)-tert-Butoxycarbonylamino-N-[(2-methoxy)ethyl]-caprolactam (71). (3S)-tert-Butoxycarbonylamino caprolactam **33** (2.0 g, 8.8 mmole) is dissolved in DMF (10 mL). Potassium *tert*-butoxide (1.25 g, 10.6 mmole) is added dropwise, and stirred for 45 min. Bromo ethyl methyl ether (0.99 mL, 10.6 mmole) is then added. The resulting mixture is stirred for 3 hr, at which time the reaction is quenched by water and extracted with ethyl acetate. The organic layer is then washed with 1N HCl, H₂O, and brine, dried over MgSO₄, filtered and evaporated to give an oil which is chromatographed over flash silica with hexanes:EtOAc (1:1) to give **71**.


(2R)-Carbo-tert-butoxymethyl decanoic acid amide of N-[(2-methoxy)ethyl]-caprolactam-(3S)-amine (73). (3S)-tert-Butoxycarbonylamino-N-[(2-methoxy)ethyl]-caprolactam **71** (1.46 g, 5.1 mmole) is dissolved in 15 mL of methylene chloride, cooled to 0°C, and trifluoroacetic acid (15 mL) is added slowly. The mixture is stirred for 2 hr at room temperature and evaporated to give **72**.

The caprolactam amine **72** is mixed with (2R)-tert-Butylcarboxymethyl decanoic acid (1.5 g, 5.2 mmole), HOBT (2.07 g, 15.3 mmole), EDAC (1.17 g, 6.1 mmole) in 15 mL of DMF and 1.43 mL (13 mmole) of NMM at 0°C, and stirred for 17 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is washed with 1N HCl, H₂O, 1N NaOH, and brine, dried over MgSO₄, filtered and evaporated to give an oil which is chromatographed over flash silica with EtOAc to give **73**.

(2R)-O-Benzyl-N-hydroxycarboxamidomethyl decanoic acid amide of N-[(2-methoxy)ethyl]-caprolactam-(3S)-amine (75). (3S)-tert-Butoxycarbonylamino-N-[(2-methoxy)ethyl]-caprolactam **73** (1.2 g, 2.64 mmole) is dissolved in 15 mL of methylene chloride, cooled to 0°C, and trifluoroacetic acid (15 mL) is added slowly. The mixture is stirred for 1.5 hr at room temperature and evaporated to give **74**.

A mixture of the crude acid **74** (1.5 g, 2.64 mmole), *O*-benzyl hydroxylamine hydrochloride (0.51 g, 3.17 mmole) and HOBT (1.07 g, 7.92 mmole) in 10 mL of DMF and 0.99 mL of NMM (9.0 mmole) is charged with EDAC (0.61 g, 3.17 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1N HCl, NaHCO₃,

36



(3S)-tert-Butoxycarbonylamino-N-[(2-methoxy)ethyl]-caprolactam (71). (3S)-tert-Butoxycarbonylamino caprolactam 33 (2.0 g, 8.8 mmole) is dissolved in DMF (10 mL). Potassium *tert*-butoxide (1.25 g, 10.6 mmole) is added dropwise, and stirred for 45 min. Bromo ethyl methyl ether (0.99 mL, 10.6 mmole) is then added. The resulting mixture is stirred for 3 hr, at which time the reaction is quenched by water and extracted with ethyl acetate. The organic layer is then washed with 1N HCl, H₂O, and brine, dried over MgSO₄, filtered and evaporated to give an oil which is chromatographed over flash silica with hexanes:EtOAc (1:1) to give 71.

(2R)-Carbo-tert-butoxymethyl decanoic acid amide of N-[(2-methoxy)ethyl]-caprolactam-(3S)-amine (73). (3S)-tert-Butoxycarbonylamino-N-[(2-methoxy)ethyl]-caprolactam 71 (1.46 g, 5.1 mmole) is dissolved in 15 mL of methylene chloride, cooled to 0°C, and trifluoroacetic acid (15 mL) is added slowly. The mixture is stirred for 2 hr at room temperature and evaporated to 72.

The caprolactam amine 72 is mixed with (2R)-tert-Butylcarboxymethyl decanoic acid (1.5 g, 5.2 mmole), HOBT (2.07 g, 15.3 mmole), EDAC (1.17 g, 6.1 mmole) in 15 mL of DMF and 1.43 mL (13 mmole) of NMM at 0°C, and stirred for 17 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is washed with 1N HCl, H₂O, 1N NaOH, and brine, dried over MgSO₄, filtered and evaporated to give an oil which is chromatographed over flash silica with EtOAc to give 73.

(2R)-O-Benzyl-N-hydroxycarboxamidomethyl decanoic acid amide of N-[(2-methoxy)ethyl]-caprolactam-(3S)-amine (75). (3S)-tert-Butoxycarbonylamino-N-[(2-methoxy)ethyl]-caprolactam 73 (1.2 g, 2.64 mmole) is dissolved in 15 mL of methylene chloride, cooled to 0°C, and trifluoroacetic acid (15 mL) is added slowly. The mixture is stirred for 1.5 hr at room temperature and evaporated to give 74.


A mixture of the crude acid 74 (1.5 g, 2.64 mmole), *O*-benzyl hydroxylamine hydrochloride (0.51 g, 3.17 mmole) and HOBT (1.07 g, 7.92 mmole) in 10 mL of DMF and 0.99 mL of NMM (9.0 mmole) is charged with EDAC (0.61 g, 3.17 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1N HCl, NaHCO₃,

and brine, dried over MgSO_4 , filtered and evaporated to give an oil which is chromatographed over flash silica with EtOAc to give 75.

(2R)-N-Hydroxycarboxamidomethyl decanoic acid amide of N-[(2-methoxy)ethyl]-caprolactam-(3S)-amine (11). The benzyl hydroxamic acid 75 (700 mg, 1.39 mmole) is taken in 10 mL of EtOH and the mixture is charged with 10% palladium on carbon (70 mg) and stirred under one atmosphere of hydrogen for 1.5 hr. The mixture is filtered through celite and concentrated to give a solid, which is purified by flash chromatography (silica, 4% formic acid in EtOAc) to give 11.

Example 12

Synthesis of (2R)-N-Hydroxycarboxamidomethyl decanoic acid amide of 1-N-n-butyl-caprolactam-(3S)-amine (12)



(3S)-tert-Butoxycarbonylamino-N-n-butyl-caprolactam (76). (3S)-tert-Butoxycarbonylamino caprolactam 33 (2.0 g, 8.8 mmole) is dissolved in DMF (8 mL). Potassium *tert*-butoxide (1.25 g, 10.6 mmole) is added dropwise, and stirred for 50 min. 1-Bromo butane (1.13 mL, 10.6 mmole) is then added. The resulting mixture is stirred for 3 hr, at which time the reaction is quenched with water and extracted with ethyl acetate. The organic layer is then washed with 1N HCl, H_2O , and brine, dried over MgSO_4 , filtered and evaporated to give an oil which is chromatographed over flash silica with hexanes:EtOAc (1:1) to give 76.


(2R)-carbo-tert-butoxymethyl decanoic acid amide of N-n-butyl-caprolactam-(3S)-amine (78). 76 (2.37g, 8.3 mmole) is dissolved in 15 mL of methylene chloride, cooled to 0°C , and trifluoroacetic acid (15 mL) is added slowly. The mixture is stirred for 2 hr at room temperature and evaporated to give 77.

and brine, dried over MgSO₄, filtered and evaporated to give an oil which is chromatographed over flash silica with EtOAc to give 75.

(2R)-N-Hydroxycarboxamidomethyl decanoic acid amide of N-[(2-methoxy)ethyl]-caprolactam-(3S)-amine (11). The benzyl hydroxamic acid 75 (700 mg, 1.39 mmole) is taken in 10 mL of EtOH and the mixture is charged with 10% palladium on carbon (70 mg) and stirred under one atmosphere of hydrogen for 1.5 hr. The mixture is filtered through celite and concentrated to give a solid, which is purified by flash chromatography (silica, 4% formic acid in EtOAc) to give 11.

Example 12

Synthesis of (2R)-N-Hydroxycarboxamidomethyl decanoic acid amide of 1-N-n-butyl-caprolactam-(3S)-amine (12)



(3S)-tert-Butoxycarbonylamino-N-n-butyl-caprolactam (76). (3S)-tert-Butoxycarbonylamino caprolactam 33 (2.0 g, 8.8 mmole) is dissolved in DMF (8 mL). Potassium *tert*-butoxide (1.25 g, 10.6 mmole) is added dropwise, and stirred for 50 min. 1-Bromo butane (1.13 mL, 10.6 mmole) is then added. The resulting mixture is stirred for 3 hr, at which time the reaction is quenched with water and extracted with ethyl acetate. The organic layer is then washed with 1N HCl, H₂O, and brine, dried over MgSO₄, filtered and evaporated to give an oil which is chromatographed over flash silica with hexanes:EtOAc (1:1) to give 76.

(2R)-carbo-tert-butoxymethyl decanoic acid amide of N-n-butyl-caprolactam-(3S)-amine (78). 76 (2.37g, 8.3 mmole) is dissolved in 15 mL of methylene chloride, cooled to 0°C, and trifluoroacetic acid (15 mL) is added slowly. The mixture is stirred for 2 hr at room temperature and evaporated to give 77.

The caprolactam amine 77 is mixed with 62 (1.5 g, 5.2 mmole), HOBT (3.36 g, 24.9 mmole), EDAC (1.92 g, 10 mmole) in 15 mL of DMF and 2.3 mL (20.8 mmole) of NMM at 0°C, and stirred for 17 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is washed with 1N HCl, H₂O, 1N NaOH, and brine, dried over MgSO₄, filtered and evaporated to give an oil which is chromatographed over flash silica with EtOAc to give 78.

(2R)-O-Benzylhydroxycarboxamidomethyl decanoic acid amide of N-n-butyl-caprolactam-(3S)-amine (80). 78 (1.4 g, 3.1 mmole) is dissolved in 15 mL of methylene chloride and cooled to 0°C and trifluoroacetic acid (15 mL) is added slowly. The mixture is stirred for 1.5 hr at room temperature and evaporated to give 79.

A mixture of the crude acid 79 (1.5 g, 3.1 mmole), O-benzyl hydroxylamine hydrochloride (0.6 g, 3.72 mmole) and HOBT (1.26 g, 9.3 mmole) in 10 mL of DMF and 1.16 mL of NMM (10.5 mmole) is charged with EDAC (0.71 g, 3.72 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is washed with 1N HCl, 1N NaOH, and brine, dried over MgSO₄, filtered and evaporated to give an oil which is chromatographed over flash silica with EtOAc to give 80.

(2R)-N-Hydroxycarboxamidomethyl decanoic acid amide of 1-N-n-butyl-caprolactam-(3S)-amine (12). The benzyl hydroxamic acid 80 (900 mg, 1.8 mmole) is taken in 10 mL of EtOH and the mixture is charged with 10% palladium on carbon (90 mg) and stirred under one atmosphere of hydrogen for 2 hr. The mixture is filtered through celite and concentrated to give a solid, which is purified by flash chromatography (silica, 2% formic acid in EtOAc). The product is then crystallized from ethyl acetate to give 12.

Example 13

(2R)-Isobutyl-(3S)-[N-hydroxycarboxamido]-butanoic acid amide of (1N)-carbomethoxymethyl-caprolactam-(3S)-amine (13)

The caprolactam amine 77 is mixed with 62 (1.5 g, 5.2 mmole), HOBT (3.36 g, 24.9 mmole), EDAC (1.92 g, 10 mmole) in 15 mL of DMF and 2.3 mL (20.8 mmole) of NMM at 0°C, and stirred for 17 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is washed with 1N HCl, H₂O, 1N NaOH, and brine, dried over MgSO₄, filtered and evaporated to give an oil which is chromatographed over flash silica with EtOAc to give 78.

(2R)-O-Benzylhydroxycarboxamidomethyl decanoic acid amide of N-n-butyl-caprolactam-(3S)-amine (80). 78 (1.4 g, 3.1 mmole) is dissolved in 15 mL of methylene chloride and cooled to 0°C and trifluoroacetic acid (15 mL) is added slowly. The mixture is stirred for 1.5 hr at room temperature and evaporated to give 79.

A mixture of the crude acid 79 (1.5 g, 3.1 mmole), O-benzyl hydroxylamine hydrochloride (0.6 g, 3.72 mmole) and HOBT (1.26 g, 9.3 mmole) in 10 mL of DMF and 1.16 mL of NMM (10.5 mmole) is charged with EDAC (0.71 g, 3.72 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is washed with 1N HCl, 1N NaOH, and brine, dried over MgSO₄, filtered and evaporated to give an oil which is chromatographed over flash silica with EtOAc to give 80.


(2R)-N-Hydroxycarboxamidomethyl decanoic acid amide of 1-N-n-butyl-caprolactam-(3S)-amine (12). The benzyl hydroxamic acid 80 (900 mg, 1.8 mmole) is taken in 10 mL of EtOH and the mixture is charged with 10% palladium on carbon (90 mg) and stirred under one atmosphere of hydrogen for 2 hr. The mixture is filtered through celite and concentrated to give a solid, which is purified by flash chromatography (silica, 2% formic acid in EtOAc). The product is then crystallized from ethyl acetate to give 12.

Example 13

(2R)-Isobutyl-(3S)-[N-hydroxycarboxamido]-butanoic acid amide of (1N)-carbomethoxymethyl-caprolactam-(3S)-amine (13)

Example 14

(2R)-Isobutyl-(3R)-[N-hydroxycarboxamido]-butanoic acid amide of (1N)-carbomethoxymethyl-caprolactam-(3S)-amine (14)



(2R)-Isobutyl-(3R,S)-carbo-*tert*-butoxybutanoic acid amide of (1N)-carbomethoxymethylcaprolactam-(3S)-amine (82). A mixture of acids **81** (3R:3S = 8:1; 720 mg, 2.95 mmole), **55** (767 mg, 3.84 mmole) and HOBT (1.39 g, 10.33 mmole) in 4 mL of DMF and 4 mL of NMM is charged with EDAC (737 mg, 3.84 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered and evaporated to give a crude solid which is chromatographed over flash silica with hexanes:EtOAc (3:1 → 1:2) to give **82** (3R:3S = 8:1).


(2R)-Isobutyl-(3R,S)-[O-benzyl-N-hydroxycarboxamido]-butanoic acid amide of (1N)-carbomethoxymethyl-caprolactam-(3S)-amine (84). The esters **82** (3R:3S = 8:1; 560 mg, 1.31 mmole), are dissolved in 4 mL of CH₂Cl₂ under argon and to this mixture is added 4 mL of trifluoroacetic acid via syringe. The mixture is stirred for 30 min. at which time the mixture is concentrated to give **83**.

A mixture of the resulting crude acid **83** (560 mg, 1.51 mmole), *O*-benzyl hydroxylamine hydrochloride (314 mg, 1.96 mmole) and HOBT (917 mg, 6.80 mmole) in 4 mL of DMF and 4 mL of NMM is charged with EDAC (376 mg, 1.96 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered and evaporated. The crude solid is filtered through a plug of silica gel and then recrystallized to give the **84** (3R) diastereomer. The mother liquor is then chromatographed through silica gel with hexanes:EtOAc (1:1 → 0:1) to give **84** (3S).

(2R)-Isobutyl-(3S)-[N-hydroxycarboxamido]-butanoic acid amide of (1N)-carbomethoxymethyl-caprolactam-(3S)-amine (13). The benzyl hydroxamic acid **84** (3S) (45 mg, 0.095 mmole) is taken in 1 mL 95% EtOH and the mixture is charged with

Example 14

(2R)-Isobutyl-(3R)-[N-hydroxycarboxamido]-butanoic acid amide of (1N)-carbomethoxymethyl-caprolactam-(3S)-amine (14)



(2R)-Isobutyl-(3R,S)-carbo-*tert*-butoxybutanoic acid amide of (1N)-carbomethoxymethylcaprolactam-(3S)-amine (82). A mixture of acids **81** (3R:3S = 8:1; 720 mg, 2.95 mmole), **55** (767 mg, 3.84 mmole) and HOBT (1.39 g, 10.33 mmole) in 4 mL of DMF and 4 mL of NMM is charged with EDAC (737 mg, 3.84 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered and evaporated to give a crude solid which is chromatographed over flash silica with hexanes:EtOAc (3:1 → 1:2) to give **82** (3R:3S = 8:1).

(2R)-Isobutyl-(3R,S)-[O-benzyl-N-hydroxycarboxamido]-butanoic acid amide of (1N)-carbomethoxymethyl-caprolactam-(3S)-amine (84). The esters **82** (3R:3S = 8:1; 560 mg, 1.31 mmole), are dissolved in 4 mL of CH₂Cl₂ under argon and to this mixture is added 4 mL of trifluoroacetic acid via syringe. The mixture is stirred for 30 min. at which time the mixture is concentrated to give **83**.

A mixture of the resulting crude acid **83** (560 mg, 1.51 mmole), O-benzyl hydroxylamine hydrochloride (314 mg, 1.96 mmole) and HOBT (917 mg, 6.80 mmole) in 4 mL of DMF and 4 mL of NMM is charged with EDAC (376 mg, 1.96 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered and evaporated. The crude solid is filtered through a plug of silica gel and then recrystallized to give the **84** (3R) diastereomer. The mother liquor is then chromatographed through silica gel with hexanes:EtOAc (1:1 → 0:1) to give **84** (3S).


(2R)-Isobutyl-(3S)-[N-hydroxycarboxamido]-butanoic acid amide of (1N)-carbomethoxymethyl-caprolactam-(3S)-amine (13). The benzyl hydroxamic acid **84** (3S) (45 mg, 0.095 mmole) is taken in 1 mL 95% EtOH and the mixture is charged with

5 mg of 10% palladium on carbon and stirred under one atmosphere of hydrogen for 12 hours. The mixture is then filtered through celite and concentrated to give a crude oil which is crystallized from EtOAc:EtOH (20:1) to give 13.

(2R)-Isobutyl-(3R)-[N-hydroxycarboxamido]-butanoic acid amide of (1N)-carbomethoxymethyl-caprolactam-(3S)-amine (14). The benzyl hydroxamic acid 84 (3R) (100 mg, 0.211 mmole) is taken in 1.5 mL 95% EtOH and the mixture is charged with 10% palladium on carbon (10 mg) and stirred under one atmosphere of hydrogen for 15 hours. The mixture is then filtered through celite and concentrated to give 14.

Example 15

Synthesis of (2R)-Isobutyl-(3S)-[N-hydroxycarboxamido]-hexanoic acid amide of (1N)-carbomethoxymethyl-caprolactam-(3S)-amine (15)



(2R)-Isobutyl-(3S)-[carbo-*tert*-butoxy]-hex-5-enoic acid amide of (1N)-carbomethoxymethylcaprolactam-(3S)-amine (86). A mixture of acid 85 (1.91 g, 7.07 mmole), 55 (1.84 g, 9.20 mmole) and HOBT (2.77 g, 20.5 mmole) in 2 mL of DMF and 2 mL of NMM is charged with EDAC (2.04 g, 10.6 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered and evaporated to give a crude solid which chromatographed over flash silica with hexanes : EtOAc (3:1 → 1:2) to give 86.


(2R)-Isobutyl-(3R,S)-carboxyhex-5-enoic acid amide of (1N)-carbomethoxymethyl-caprolactam-(3S)-amine (87). Trifluoroacetic acid (15 mL) is added via syringe to a solution of *tert*-butyl esters 86 (3R:3S = 2:3; 1.51g, 3.34 mmole) in 15 mL of CH₂Cl₂ under argon and the resulting mixture is stirred for 1 hr at room

5 mg of 10% palladium on carbon and stirred under one atmosphere of hydrogen for 12 hours. The mixture is then filtered through celite and concentrated to give a crude oil which is crystallized from EtOAc:EtOH (20:1) to give 13.

(2R)-Isobutyl-(3R)-[N-hydroxycarboxamido]-butanoic acid amide of (1N)-carbomethoxymethyl-caprolactam-(3S)-amine (14). The benzyl hydroxamic acid **84** (3R) (100 mg, 0.211 mmole) is taken in 1.5 mL 95% EtOH and the mixture is charged with 10% palladium on carbon (10 mg) and stirred under one atmosphere of hydrogen for 15 hours. The mixture is then filtered through celite and concentrated to give 14.

Example 15

Synthesis of (2R)-Isobutyl-(3S)-[N-hydroxycarboxamido]-hexanoic acid amide of (1N)-carbomethoxymethyl-caprolactam-(3S)-amine (15)



(2R)-Isobutyl-(3S)-[carbo-*tert*-butoxy]-hex-5-enoic acid amide of (1N)-carbomethoxymethylcaprolactam-(3S)-amine (86). A mixture of acid **85** (1.91 g, 7.07 mmole), **55** (1.84 g, 9.20 mmole) and HOBt (2.77 g, 20.5 mmole) in 2 mL of DMF and 2 mL of NMM is charged with EDAC (2.04 g, 10.6 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered and evaporated to give a crude solid which chromatographed over flash silica with hexanes : EtOAc (3:1 → 1:2) to give **86**.

(2R)-Isobutyl-(3R,S)-carboxyhex-5-enoic acid amide of (1N)-carbomethoxymethyl-caprolactam-(3S)-amine (87). Trifluoroacetic acid (15 mL) is added via syringe to a solution of *tert*-butyl esters **86** (3R:3S = 2:3; 1.51g, 3.34 mmole) in 15 mL of CH₂Cl₂ under argon and the resulting mixture is stirred for 1 hr at room


temperature. The material is then concentrated under vacuum to give **87** as a diastereomeric mixture (3R:3S = 2:3) which is carried forward without purification.

(2R)-Isobutyl-(3S)-[O-benzyl-N-hydroxycarboxamido]-hex-5-enoic acid amide of (1N)-carbomethoxymethyl-caprolactam-(3S)-amine (**88**). A mixture of acids **87** (3R:3S = 2:3; 1.32 g, 3.34 mmole), O-benzyl hydroxylamine hydrochloride (0.693 g, 4.34 mmole) and HOBt (1.26 g, 9.35 mmole) in 7 mL of DMF and 7 mL of NMM is charged with EDAC (0.962 mg, 5.01 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, and evaporated to give a crude solid which is filtered through a plug of silica gel with EtOAc and then recrystallized 2x from hexanes:EtOAc (2:1) to give **88**.

(2R)-Isobutyl-(3S)-[N-hydroxycarboxamido]-hexanoic acid amide of (1N)-carbomethoxymethyl-caprolactam-(3S)-amine (**15**). The benzyl hydroxamic acid **88** (176 mg, 0.351 mmole) is taken in 1 mL of 95% EtOH and the mixture is charged with 30 mg of 10% palladium on carbon and stirred under one atmosphere of hydrogen for 12 hours. The mixture is then filtered through celite and concentrated to give **15**.

Example 16

(2R)-Isobutyl-(3S)-[N-hydroxycarboxamido]-6-hydroxyhexanoic acid amide of (1N)-carbomethoxymethyl-caprolactam-(3S)-amine (**16**)



(2R)-Isobutyl-(3S)-[O-benzyl-N-hydroxycarboxamido]-6-hydroxyhexanoic acid amide of (1N)-carbomethoxymethyl-caprolactam-(3S)-amine (**89**). The starting olefin **88** (3R:3S = ~1:1; 240 mg, 0.479 mmol) is taken in 5 mL of dry THF under argon and cooled to 0°C at which time 9-borabicyclononane ("9-BBN") (3.45 mL, 0.5M in THF,


temperature. The material is then concentrated under vacuum to give **87** as a diastereomeric mixture (3R:3S = 2:3) which is carried forward without purification.

(2R)-Isobutyl-(3S)-[O-benzyl-N-hydroxycarboxamido]-hex-5-enoic acid amide of (1N)-carbomethoxymethyl-caprolactam-(3S)-amine (**88**). A mixture of acids **87** (3R:3S = 2:3; 1.32 g, 3.34 mmole), O-benzyl hydroxylamine hydrochloride (0.693 g, 4.34 mmole) and HOBT (1.26 g, 9.35 mmole) in 7 mL of DMF and 7 mL of NMM is charged with EDAC (0.962 mg, 5.01 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, and evaporated to give a crude solid which is filtered through a plug of silica gel with EtOAc and then recrystallized 2x from hexanes:EtOAc (2:1) to give **88**.

(2R)-Isobutyl-(3S)-[N-hydroxycarboxamido]-hexanoic acid amide of (1N)-carbomethoxymethyl-caprolactam-(3S)-amine (**15**). The benzyl hydroxamic acid **88** (176 mg, 0.351 mmole) is taken in 1 mL of 95% EtOH and the mixture is charged with 30 mg of 10% palladium on carbon and stirred under one atmosphere of hydrogen for 12 hours. The mixture is then filtered through celite and concentrated to give **15**.

Example 16

(2R)-Isobutyl-(3S)-[N-hydroxycarboxamido]-6-hydroxyhexanoic acid amide of (1N)-carbomethoxymethyl-caprolactam-(3S)-amine (**16**)



(2R)-Isobutyl-(3S)-[O-benzyl-N-hydroxycarboxamido]-6-hydroxyhexanoic acid amide of (1N)-carbomethoxymethyl-caprolactam-(3S)-amine (**89**). The starting olefin **88** (3R:3S = ~1:1; 240 mg, 0.479 mmol) is taken in 5 mL of dry THF under argon and cooled to 0°C at which time 9-borabicyclononane ("9-BBN") (3.45 mL, 0.5M in THF,

1.73 mmol) is added via syringe and the solution is stirred at 0°C for 2 hr at which time NaOH (2 mL, 1M) and H₂O₂ (2 mL, 30%) are added and the resulting solution stirred for 5 min. The mixture is then partitioned between EtOAc and water and the organic layer is washed with sat. NH₄Cl and brine, dried over MgSO₄, filtered and evaporated to give a crude oil. This diastereomeric mixture is chromatographed 2x over flash silica with EtOAc:EtOH (20:1) to give the 1S-89.


(2R)-Isobutyl-(3S)-[N-hydroxycarboxamido]-6-hydroxyhexanoic acid amide of (1N)-carbomethoxymethyl-caprolactam-(3S)-amine (16). The benzyl hydroxamic acid 89 (45 mg, 0.351 mmole) is taken in 1 mL of 95% EtOH and the mixture is charged with 10 mg of 10% palladium on carbon and stirred under one atmosphere of hydrogen for 12 hours. The mixture is then filtered through celite and concentrated to give 16.

Example 17

Synthesis of (2R)-[(1S)-N-hydroxycarboxamido]-ethyldecanoic acid amide of 1N-(carbomethoxy-methyl)-caprolactam-(3S)-amine (17)

Example 18

Synthesis of (2R)-[(1R)-N-hydroxycarboxamido]-ethyldecanoic acid amide of 1N-(carbomethoxy-methyl)-caprolactam-(3S)-amine (18)



1.73 mmol) is added via syringe and the solution is stirred at 0°C for 2 hr at which time NaOH (2 mL, 1M) and H₂O₂ (2 mL, 30%) are added and the resulting solution stirred for 5 min. The mixture is then partitioned between EtOAc and water and the organic layer is washed with sat. NH₄Cl and brine, dried over MgSO₄, filtered and evaporated to give a crude oil. This diastereomeric mixture is chromatographed 2x over flash silica with EtOAc:EtOH (20:1) to give the 1S-89.


(2R)-Isobutyl-(3S)-[N-hydroxycarboxamido]-6-hydroxyhexanoic acid amide of (1N)-carbomethoxymethyl-caprolactam-(3S)-amine (16). The benzyl hydroxamic acid 89 (45 mg, 0.351 mmole) is taken in 1 mL of 95% EtOH and the mixture is charged with 10 mg of 10% palladium on carbon and stirred under one atmosphere of hydrogen for 12 hours. The mixture is then filtered through celite and concentrated to give 16.

Example 17

Synthesis of (2R)-[(1S)-N-hydroxycarboxamido]-ethyldecanoic acid amide of 1N-(carbomethoxy-methyl)-caprolactam-(3S)-amine (17)

Example 18

Synthesis of (2R)-[(1R)-N-hydroxycarboxamido]-ethyldecanoic acid amide of 1N-(carbomethoxy-methyl)-caprolactam-(3S)-amine (18)



(2R)-[(1R)-Carbo-*tert*-Butoxy]-ethyl decanoic acid (90). The starting acid 62 (4.5 g, 15.73 mmole) is dissolved in 5 mL of dry THF under an argon atmosphere and cooled to -78°C at which time lithium hexamethyldisilazane (39.3 mL, 1 M in THF, 39.3 mmole) is added via syringe and the mixture is stirred for 20 min. The mixture is allowed to warm to 0°C and then re-cooled to -78°C. Methyl iodide (1.08 mL, 17.3 mmole) is then added slowly via syringe and the resulting mixture is stirred for 2 hr at -78°C and 30 min. at 0°C, at which time the reaction is quenched with saturated NH₄Cl, partitioned between EtOAc and water and layers separated. The organic layer is washed with NH₄Cl, and brine, dried over MgSO₄, filtered and evaporated to give 5.81 g of crude oil which is chromatographed over flash silica with hexane:EtOAc (2:1) to give 90.

(2R)-[(1R)-Carbo-*tert*-Butoxy]-ethyl decanoic acid (91). The predominately (1R) acid 90 (1R:1S = 8:1, 5 g, 15.7 mmole) is dissolved in 5 mL of dry THF under an argon atmosphere and cooled to -78°C. At which time lithium diisopropylamine (47 mL, 2 M in THF, 47 mmole) is added via syringe and the mixture is stirred for 16 hr at -78°C to 22°C as the dry ice-acetone bath warmed to room temperature. The mixture is then re-cooled to -78°C, quenched with 15 mL of methanol (78 mmole) and poured into NH₄Cl. After warming up to room temperature, the reaction mixture is partitioned between EtOAc and water and layers separated. The organic layer is washed with 1 N HCl, and brine, dried over MgSO₄, filtered and evaporated to give 5.81 g of crude oil which is chromatographed over flash silica with hexane:EtOAc (2:1) to give 91 (1S:1R = 2:3).

(2R)-[(1R,S)-Carbo-*tert*-butoxy]-ethyldecanoic acid amide of 1N-(carbomethoxy-methyl)-caprolactam-(3S)-amine (92). A mixture of acids 91 (1R:1S = 3:2, 1.5 g, 5.00 mmole), the amine 55 (1.5 g, 7.50 mmole) and HOBt (1.53 g, 10.00 mmole) in 10 mL of DMF and 1.1 mL of NMM is charged with EDAC (1.15 mg, 6.00 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered and evaporated to a crude solid which is chromatographed over flash silica with hexanes:EtOAc (1:1) to give 92 (1R:1S = 3:2).

(2R)-[(1R)-O-Benzyl-N-hydroxycarboxamido]-ethyldecanoic acid amide of 1N-(carbomethoxymethyl)-caprolactam-(3S)-amine (94). (2R)-[(1S)-O-Benzyl-N-hydroxycarboxamido]-ethyldecanoic acid amide of 1N-(carbomethoxymethyl)-caprolactam-(3S)-amine (95). The esters 92 (1R:1S = 3:2, 1.40 g, 2.90 mmole), are dissolved in 7 mL of CH₂Cl₂ under argon and to this mixture is added 7 mL of trifluoroacetic acid via

(2R)-[(1R)-Carbo-*tert*-Butoxy]-ethyl decanoic acid (90). The starting acid **62** (4.5 g, 15.73 mmole) is dissolved in 5 mL of dry THF under an argon atmosphere and cooled to -78°C at which time lithium hexamethyldisilazane (39.3 mL, 1 M in THF, 39.3 mmole) is added via syringe and the mixture is stirred for 20 min. The mixture is allowed to warm to 0°C and then recooled to -78°C. Methyl iodide (1.08 mL, 17.3 mmole) is then added slowly via syringe and the resulting mixture is stirred for 2 hr at -78°C and 30 min. at 0°C, at which time the reaction is quenched with saturated NH₄Cl, partitioned between EtOAc and water and layers separated. The organic layer is washed with NH₄Cl, and brine, dried over MgSO₄, filtered and evaporated to give 5.81 g of crude oil which is chromatographed over flash silica with hexane:EtOAc (2:1) to give **90**.

(2R)-[(1R)-Carbo-*tert*-Butoxy]-ethyl decanoic acid (91). The predominately (1R) acid **90** (1R:1S = 8:1, 5 g, 15.7 mmole) is dissolved in 5 mL of dry THF under an argon atmosphere and cooled to -78°C. At which time lithium diisopropylamine (47 mL, 2 M in THF, 47 mmole) is added via syringe and the mixture is stirred for 16 hr at -78°C to 22°C as the dry ice-acetone bath warmed to room temperature. The mixture is then recooled to -78°C, quenched with 15 mL of methanol (78 mmole) and poured into NH₄Cl. After warming up to room temperature, the reaction mixture is partitioned between EtOAc and water and layers separated. The organic layer is washed with 1 N HCl, and brine, dried over MgSO₄, filtered and evaporated to give 5.81 g of crude oil which is chromatographed over flash silica with hexane:EtOAc (2:1) to give **91** (1S:1R = 2:3).

(2R)-[(1R,S)-Carbo-*tert*-butoxy]-ethyldecanoic acid amide of 1N-(carbomethoxy-methyl)-caprolactam-(3S)-amine (92). A mixture of acids **91** (1R:1S = 3:2, 1.5 g, 5.00 mmole), the amine **55** (1.5 g, 7.50 mmole) and HOBT (1.53 g, 10.00 mmole) in 10 mL of DMF and 1.1 mL of NMM is charged with EDAC (1.15 mg, 6.00 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered and evaporated to a crude solid which is chromatographed over flash silica with hexanes:EtOAc (1:1) to give **92** (1R:1S = 3:2).

(2R)-[(1R)-O-Benzyl-N-hydroxycarboxamido]-ethyldecanoic acid amide of 1N-(carbomethoxymethyl)-caprolactam-(3S)-amine (94). (2R)-[(1S)-O-Benzyl-N-hydroxycarboxamido]-ethyldecanoic acid amide of 1N-(carbomethoxymethyl)-caprolactam-(3S)-amine (**95**). The esters **92** (1R:1S = 3:2, 1.40 g, 2.90 mmole), are dissolved in 7 mL of CH₂Cl₂ under argon and to this mixture is added 7 mL of trifluoroacetic acid via

syringe. The mixture is stirred for 2 hr at which time the mixture is concentrated to give **93**.

A mixture of the resulting crude acid **93** (1.27 g, 2.86 mmole), *O*-benzyl hydroxylamine hydrochloride (456 mg, 2.86 mmole) and HOBT (875 mg, 5.72 mmole) in 5 mL of DMF and 941 μ L of NMM is charged with EDAC (658 mg, 3.43 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered and evaporated. The crude solid is then chromatographed through silica gel with hexanes:EtOAc (1:2) to give **94** and **95**.

(2R)-[(1S)-*N*-hydroxycarboxamido]-ethyldecanoic acid amide of 1*N*-(carbomethoxy-methyl)-caprolactam-(3S)-amine (**17**). The benzyl hydroxamic acid **95** (260 mg, 0.489 mmole) is taken in 6 mL EtOH and the mixture is charged with 10% palladium on carbon (60 mg) and stirred under one atmosphere of hydrogen for 45 min. The mixture is then filtered through celite, concentrated and then recrystallized from EtOAc to give **17**.

(2R)-[(1R)-*N*-hydroxycarboxamido]-ethyldecanoic acid amide of 1*N*-(carbomethoxy-methyl)-caprolactam-(3S)-amine (**18**). The benzyl hydroxamic acid **94** (300 mg, 0.565 mmole) is taken in 6 mL of EtOH and the mixture is charged with 10% palladium on carbon (60 mg) and stirred under one atmosphere of hydrogen for 45 min. The mixture is then filtered through celite, concentrated and then recrystallized from EtOAc to give **18**.

Example 19

Synthesis of (2R)-[(1S)-*N*-Hydroxycarboxamido]-ethyl decanoic acid amide of N-[(2-methoxy)ethyl]-caprolactam-(3S)-amine (**19**)

syringe. The mixture is stirred for 2 hr at which time the mixture is concentrated to give 93.


A mixture of the resulting crude acid 93 (1.27 g, 2.86 mmole), *O*-benzyl hydroxylamine hydrochloride (456 mg, 2.86 mmole) and HOBT (875 mg, 5.72 mmole) in 5 mL of DMF and 941 μ L of NMM is charged with EDAC (658 mg, 3.43 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered and evaporated. The crude solid is then chromatographed through silica gel with hexanes:EtOAc (1:2) to give 94 and 95.

(2R)-[(1S)-N-hydroxycarboxamido]-ethyldecanoic acid amide of 1N-(carbomethoxy-methyl)-caprolactam-(3S)-amine (17). The benzyl hydroxamic acid 95 (260 mg, 0.489 mmole) is taken in 6 mL EtOH and the mixture is charged with 10% palladium on carbon (60 mg) and stirred under one atmosphere of hydrogen for 45 min. The mixture is then filtered through celite, concentrated and then recrystallized from EtOAc to give 17.

(2R)-[(1R)-N-hydroxycarboxamido]-ethyldecanoic acid amide of 1N-(carbomethoxy-methyl)-caprolactam-(3S)-amine (18). The benzyl hydroxamic acid 94 (300 mg, 0.565 mmole) is taken in 6 mL of EtOH and the mixture is charged with 10% palladium on carbon (60 mg) and stirred under one atmosphere of hydrogen for 45 min. The mixture is then filtered through celite, concentrated and then recrystallized from EtOAc to give 18.


Example 19

Synthesis of (2R)-[(1S)-N-Hydroxycarboxamido]-ethyl decanoic acid amide of N-[(2-methoxy)ethyl]-caprolactam-(3S)-amine (19)

Example 20Synthesis of (2R)-[(1R)-N-Hydroxycarboxamido]-ethyl decanoic acid amide of N-[(2-methoxy)ethyl]-caprolactam-(3S)-amine (20)

(2R)-[(1R,S)-Tert-Butylcarboxy]-ethyl decanoic acid amide of N-[(2-methoxy)ethyl]-caprolactam-(3S)-amine (96). (3S)-Amino-N-[(2-methoxy)ethyl]-caprolactam 72 (2.0 g, 10.78 mmole) is mixed with the decanoic acid 91 (1.5 g, 5.2 mmole), HOBT (3.65 g, 27 mmole), EDAC (2.07 g, 10.8 mmole) in 15 mL of DMF and 3 mL (27 mmole) of NMM at 0°C, and stirred for 17 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is washed with 1N HCl, H₂O, 1N NaOH, and brine, dried over MgSO₄, filtered and evaporated to give an oil which is chromatographed over flash silica with hexane:EtOAc (1:1) to give 96.

(2R)-[(1S)-O-Benzylhydroxycarboxamic]-ethyl decanoic acid amide of N-[(2-methoxy)ethyl]-caprolactam-(3S)-amine (98). (2R)-[(1S)-tert-butyl carboxy]-ethyl decanoic acid amide of N-[(2-methoxy)ethyl]-caprolactam-(3S)-amine 96 (1.95 g, 4.1 mmole) is dissolved in 20 mL of methylene chloride and cooled to 0°C, trifluoroacetic acid (20 mL) is added slowly. The mixture is stirred for 2 hr at room temperature and evaporated to give 97. A mixture of the crude acid 97 (2.3 g, 4.1 mmole), O-benzyl hydroxylamine hydrochloride (0.78 g, 4.92 mmole) and HOBT (1.66 g, 12.3 mmole) in 15 mL of DMF and 1.54 mL of NMM (14 mmole) is charged with EDAC (0.94 g, 4.92 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is washed with 1N HCl, 1N NaOH, and brine, dried over MgSO₄, filtered and evaporated to give an oil which is chromatographed over flash silica with EtOAc:CH₂Cl₂:CH₃OH = 10:10:1 to give the

Example 20Synthesis of (2R)-[(1R)-N-Hydroxycarboxamido]-ethyl decanoic acid amide of N-[(2-methoxy)ethyl]-caprolactam-(3S)-amine (20)

(2R)-[(1R,S)-*Tert*-Butylcarboxy]-ethyl decanoic acid amide of N-[(2-methoxy)ethyl]-caprolactam-(3S)-amine (96). (3S)-Amino-N-[(2-methoxy)ethyl]-caprolactam 72 (2.0 g, 10.78 mmole) is mixed with the decanoic acid 91 (1.5 g, 5.2 mmole), HOBT (3.65 g, 27 mmole), EDAC (2.07 g, 10.8 mmole) in 15 mL of DMF and 3 mL (27 mmole) of NMM at 0°C, and stirred for 17 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is washed with 1N HCl, H₂O, 1N NaOH, and brine, dried over MgSO₄, filtered and evaporated to give an oil which is chromatographed over flash silica with hexane:EtOAc (1:1) to give 96.

(2R)-[(1S)-O-Benzylhydroxycarboxamic]-ethyl decanoic acid amide of N-[(2-methoxy)ethyl]-caprolactam-(3S)-amine (98). (2R)-[(1S)-*tert*-butyl carboxy]-ethyl decanoic acid amide of N-[(2-methoxy)ethyl]-caprolactam-(3S)-amine 96 (1.95 g, 4.1 mmole) is dissolved in 20 mL of methylene chloride and cooled to 0°C, trifluoroacetic acid (20 mL) is added slowly. The mixture is stirred for 2 hr at room temperature and evaporated to give 97. A mixture of the crude acid 97 (2.3 g, 4.1 mmole), O-benzyl hydroxylamine hydrochloride (0.78 g, 4.92 mmole) and HOBT (1.66 g, 12.3 mmole) in 15 mL of DMF and 1.54 mL of NMM (14 mmole) is charged with EDAC (0.94 g, 4.92 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is washed with 1N HCl, 1N NaOH, and brine, dried over MgSO₄, filtered and evaporated to give an oil which is chromatographed over flash silica with EtOAc:CH₂Cl₂:CH₃OH = 10:10:1) to give the

desired product as a diastereomeric mixture (1R:1S = 3:2). Separation of single isomers is achieved by crystallization from EtOAc:hexane (3:2) to give 1S-98.


(2R)-[(1R)-O-Benzylhydroxycarboxamido]-ethyl decanoic acid amide of N-[(2-methoxy)ethyl]-caprolactam-(3S)-amine (99). The mother liquor is concentrated to give 1R-99.

(2R)-[(1S)-N-Hydroxycarboxamido]-ethyl decanoic acid amide of N-[(2-methoxy)ethyl]-caprolactam-(3S)-amine (19). The benzyl hydroxamic acid **98** (280 mg, 0.54 mmole) is taken in 8 mL of EtOH and the mixture is charged with 10% palladium on carbon (28 mg) and stirred under one atmosphere of hydrogen for 2 hr. The mixture is filtered through celite and concentrated to give a solid, which is purified by crystallization from EtOAc:CH₃OH (10:1) to give **19**.

(2R)-[(1R)-N-Hydroxycarboxamido]-ethyl decanoic acid amide of N-[(2-methoxy)ethyl]-caprolactam-(3S)-amine (20). The benzyl hydroxamic acid **99** (670 mg, 1.3 mmole) is taken in 10 mL of EtOH and the mixture is charged with 10% palladium on carbon (66 mg) and stirred under one atmosphere of hydrogen for 3.5 hr. The mixture is filtered through celite and concentrated to give a solid, which is purified by crystallization from EtOAc to give **20**.

Example 21

Synthesis of (2R)-Isobutyl-3-(N-hydroxycarboxamido)-propanoic acid amide of N-(carbomethoxymethyl)-valerolactam-(3S)-amine



(3S)-Amino-valerolactam (101). To a 1 L 3 neck round bottom flask equipped with a condenser, thermometer, magnetic stirrer and argon inlet, is added L-ornithine hydrochloride **100** (15 g, 89 mmole), acetonitrile (280 mL) and hexamethyldisilazane (100 mL, 620 mmole). The mixture is heated to reflux for two days, then cooled to

desired product as a diastereomeric mixture (1R:1S = 3:2). Separation of single isomers is achieved by crystallization from EtOAc:hexane (3:2) to give 1S-98.


(2R)-[(1R)-O-Benzylhydroxycarboxamic]-ethyl decanoic acid amide of N-[(2-methoxy)ethyl]-caprolactam-(3S)-amine (99). The mother liquor is concentrated to give 1R-99.

(2R)-[(1S)-N-Hydroxycarboxamido]-ethyl decanoic acid amide of N-[(2-methoxy)ethyl]-caprolactam-(3S)-amine (19). The benzyl hydroxamic acid 98 (280 mg, 0.54 mmole) is taken in 8 mL of EtOH and the mixture is charged with 10% palladium on carbon (28 mg) and stirred under one atmosphere of hydrogen for 2 hr. The mixture is filtered through celite and concentrated to give a solid, which is purified by crystallization from EtOAc:CH₃OH (10:1) to give 19.

(2R)-[(1R)-N-Hydroxycarboxamido]-ethyl decanoic acid amide of N-[(2-methoxy)ethyl]-caprolactam-(3S)-amine (20). The benzyl hydroxamic acid 99 (670 mg, 1.3 mmole) is taken in 10 mL of EtOH and the mixture is charged with 10% palladium on carbon (66 mg) and stirred under one atmosphere of hydrogen for 3.5 hr. The mixture is filtered through celite and concentrated to give a solid, which is purified by crystallization from EtOAc to give 20.

Example 21

Synthesis of (2R)-Isobutyl-3-(N-hydroxycarboxamido)-propanoic acid amide of N-(carbomethoxymethyl)-valerolactam-(3S)-amine



(3S)-Amino-valerolactam (101). To a 1 L 3 neck round bottom flask equipped with a condenser, thermometer, magnetic stirrer and argon inlet, is added L-ornithine hydrochloride 100 (15 g, 89 mmole), acetonitrile (280 mL) and hexamethyldisilazane (100 mL, 620 mmole). The mixture is heated to reflux for two days, then cooled to

room temperature and poured into 500 mL of methanol. The solvents are removed by rotary evaporation. The residue is taken up in 250 mL of methylene chloride, then treated with activated carbon, filtered and concentrated to a solid **101**. The compound is used without purification.

(3S)-tert-Butoxycarbonylamino valerolactam (102). (3S)-Amino-valerolactam (crude, 10.1 g, 88.4 mmole) is mixed with di-*tert*-butyl-dicarbonate **101** (19.9 g, 88.4 mmole) in 50 mL of methyl sulfoxide. The mixture is stirred overnight. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 0.1N HCl, H₂O, and brine, dried over MgSO₄, filtered and evaporated to give an oil which is chromatographed over flash silica with hexanes:EtOAc (1:1) to give **102**.

(3S)-tert-Butoxycarbonylamino-N-carbomethoxymethyl-valerolactam (103). (3S)-*tert*-Butoxycarbonylamino valerolactam **102** (1.85 g, 8.63 mmole) is dissolved in THF (5 mL) and cooled to -78°C. Lithium bis(trimethylsilyl)amide (11 mL, 10.4 mmole, 1 M in THF) is added dropwise. After 10 min, methyl bromoacetate (1.1 mL, 11.2 mmole) is added. The resulting mixture is stirred at -78°C for 2 hr, and at room temperature for 1 hr. The reaction is quenched with NH₄Cl, extracted by ethyl acetate. The organic layer is then washed with 0.1N HCl, H₂O, and brine, dried over MgSO₄, filtered and evaporated to give an oil which is chromatographed over flash silica with hexanes:EtOAc (1:1) to give **103**.

[(2R)-Isobutyl-3-carbo-*tert*-butoxy]-propanoic acid amide of N-carbomethoxymethyl-valerolactam-(3S)-amine (105). (3S)-*tert*-Butoxycarbonylamino-N-carbomethoxymethyl-valerolactam **103** (2.1 g, 7.3 mmole) is dissolved in 20 mL of methylene chloride, trifluoroacetic acid (20 mL) is added slowly. The mixture is stirred for 2 hr and evaporated to give **104**. The valerolactam amine **104** is mixed with (2R)-isobutyl-3-carbo-*tert*-butoxy-propanoic acid **23** (2.52g, 11 mmole), HOBT (3.0 g, 21.9 mmole), EDAC (2.1 g, 11 mmole) in 10 mL of DMF and 4.2 mL (29.2 mmole) of triethylamine at 0°C, and stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered and evaporated to give an oil which is chromatographed over flash silica with EtOAc to give **105**.

[(2R)-Isobutyl-3-(O-benzyl-N-hydroxycarboxamido)]-propanoic acid amide of N-(carbomethoxymethyl)-valerolactam-(3S)-amine (107). [(2R)-Isobutyl-3-*tert*-butyl-carboxy]-propanoic acid amide of N-carbomethoxymethyl-valerolactam-(3S)-amine **105** (2.2 g, 5.5 mmole) is dissolved in 10 mL of methylene chloride and trifluoroacetic acid (10 mL) is added slowly. The mixture is stirred for 2 hr and evaporated to give **106**. A mixture of the crude acid **106** (2.8 g, 5.5 mmole), O-benzyl hydroxylamine hydrochloride (1.96 g, 12.3 mmole) and HOBT (3.3 g, 24.6 mmole) in 10 mL of DMF

room temperature and poured into 500 mL of methanol. The solvents are removed by rotary evaporation. The residue is taken up in 250 mL of methylene chloride, then treated with activated carbon, filtered and concentrated to a solid 101. The compound is used without purification.

(3S)-tert-Butoxycarbonylamino valerolactam (102). (3S)-Amino-valerolactam (crude, 10.1 g, 88.4 mmole) is mixed with di-*tert*-butyl-dicarbonate 101 (19.9 g, 88.4 mmole) in 50 mL of methyl sulfoxide. The mixture is stirred overnight. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 0.1N HCl, H₂O, and brine, dried over MgSO₄, filtered and evaporated to give an oil which is chromatographed over flash silica with hexanes:EtOAc (1:1) to give 102.

3S)-tert-Butoxycarbonylamino-N-carbomethoxymethyl-valerolactam (103). (3S)-*tert*-Butoxycarbonylamino valerolactam 102 (1.85 g, 8.63 mmole) is dissolved in THF (5 mL) and cooled to -78°C. Lithium bis(trimethylsilyl)amide (11 mL, 10.4 mmole, 1 M in THF) is added dropwise. After 10 min, methyl bromoacetate (1.1 mL, 11.2 mmole) is added. The resulting mixture is stirred at -78°C for 2 hr, and at room temperature for 1 hr. The reaction is quenched with NH₄Cl, extracted by ethyl acetate. The organic layer is then washed with 0.1N HCl, H₂O, and brine, dried over MgSO₄, filtered and evaporated to give an oil which is chromatographed over flash silica with hexanes:EtOAc (1:1) to give 103.

[(2R)-Isobutyl-3-carbo-*tert*-butoxy]-propanoic acid amide of N-carbomethoxymethyl-valerolactam-(3S)-amine (105). (3S)-*tert*-Butoxycarbonylamino-N-carbomethoxymethyl-valerolactam 103 (2.1 g, 7.3 mmole) is dissolved in 20 mL of methylene chloride, trifluoroacetic acid (20 mL) is added slowly. The mixture is stirred for 2 hr and evaporated to give 104. The valerolactam amine 104 is mixed with (2R)-isobutyl-3-carbo-*tert*-butoxy-propanoic acid 23 (2.52g, 11 mmole), HOBT (3.0 g, 21.9 mmole), EDAC (2.1 g, 11 mmole) in 10 mL of DMF and 4.2 mL (29.2 mmole) of triethylamine at 0°C, and stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered and evaporated to give an oil which is chromatographed over flash silica with EtOAc to give 105.


[(2R)-Isobutyl-3-(O-benzyl-N-hydroxycarboxamido)]-propanoic acid amide of N-(carbomethoxymethyl)-valerolactam-(3S)-amine (107). [(2R)-Isobutyl-3-*tert*-butyl-carboxy]-propanoic acid amide of N-carbomethoxymethyl-valerolactam-(3S)-amine 105 (2.2 g, 5.5 mmole) is dissolved in 10 mL of methylene chloride and trifluoroacetic acid (10 mL) is added slowly. The mixture is stirred for 2 hr and evaporated to give 106. A mixture of the crude acid 106 (2.8 g, 5.5 mmole), O-benzyl hydroxylamine hydrochloride (1.96 g, 12.3 mmole) and HOBT (3.3 g, 24.6 mmole) in 10 mL of DMF

and 2 mL of NMM (28.7 mmole) is charged with EDAC (2.0 g, 10.7 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered and evaporated to give an oil which chromatographed over flash silica with hexanes:EtOAc (1:1 → 1:4) to give 107.

(2R)-Isobutyl-3-(N-hydroxycarboxamido)-propanoic acid amide of N-(carbomethoxymethyl)-valerolactam-(3S)-amine (21). The benzyl hydroxamic acid 107 (850 mg, 1.90 mmole) is taken in 10 mL EtOH and the mixture is charged with 10% palladium on carbon (85 mg) and stirred under one atmosphere of hydrogen for 1 hr. The mixture is then filtered through celite and concentrated to an oil. The crude product is purified on a silica gel column using 2% formic acid in ethyl acetate as the eluent to give 21.

Example 22

Synthesis of [(2R)-Isobutyl-3-(N-hydroxylcarboxamido)]-propanoic acid amide of 2-oxo-3-amino-N-(carbomethoxymethyl)-pyridinone (22)



2-Oxo-3-nitro-N-(carbomethoxymethyl)-pyridinone (109). 2-Hydroxy-3-nitropyridine 108 (10 g, 71.38 mmole) is mixed with pulverized potassium carbonate (10.9 g, 78.5 mmole) and 30 mL of DMF. After 10 min, methyl bromoacetate (10.4 mL, 107 mmole) is added. The resulting mixture is stirred at room temperature for 4.5 hr. The reaction is quenched by water and extracted by ethyl acetate. The organic layer is then washed with brine, dried over MgSO₄, filtered and evaporated to give an oil which is chromatographed over flash silica with EtOAc to give 109.


2-Oxo-3-amino-N-(carbomethoxymethyl)-pyridinone (110). The pyridinone 109 (3.0 g, 14.1 mmole) is taken in 45 mL of EtOH and the mixture is charged with 10% palladium on carbon (0.3 g) and stirred under one atmosphere of hydrogen for 17 hr. The mixture is then filtered through celite and concentrated to an oil. The crude product is chromatographed over flash silica with hexane:ethyl acetate (1:1 to 100% EtOAc) to give 110.

and 2 mL of NMM (28.7 mmole) is charged with EDAC (2.0 g, 10.7 mmole) and the reaction is stirred for 15 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered and evaporated to give an oil which chromatographed over flash silica with hexanes:EtOAc (1:1 → 1:4) to give 107.

(2R)-Isobutyl-3-(N-hydroxycarboxamido)-propanoic acid amide of N-(carbomethoxymethyl)-valerolactam-(3S)-amine (21). The benzyl hydroxamic acid 107 (850 mg, 1.90 mmole) is taken in 10 mL EtOH and the mixture is charged with 10% palladium on carbon (85 mg) and stirred under one atmosphere of hydrogen for 1 hr. The mixture is then filtered through celite and concentrated to an oil. The crude product is purified on a silica gel column using 2% formic acid in ethyl acetate as the eluent to give 21.

Example 22

Synthesis of [(2R)-Isobutyl-3-(N-hydroxylcarboxamido)]-propanoic acid amide of 2-oxo-3-amino-N-(carbomethoxymethyl)-pyridinone (22)



2-Oxo-3-nitro-N-(carbomethoxymethyl)-pyridinone (109). 2-Hydroxy-3-nitropyridine 108 (10 g, 71.38 mmole) is mixed with pulverized potassium carbonate (10.9 g, 78.5 mmole) and 30 mL of DMF. After 10 min, methyl bromoacetate (10.4 mL, 107 mmole) is added. The resulting mixture is stirred at room temperature for 4.5 hr. The reaction is quenched by water and extracted by ethyl acetate. The organic layer is then washed with brine, dried over MgSO₄, filtered and evaporated to give an oil which is chromatographed over flash silica with EtOAc to give 109.

2-Oxo-3-amino-N-(carbomethoxymethyl)-pyridinone (110). The pyridinone 109 (3.0 g, 14.1 mmole) is taken in 45 mL of EtOH and the mixture is charged with 10% palladium on carbon (0.3 g) and stirred under one atmosphere of hydrogen for 17 hr. The mixture is then filtered through celite and concentrated to an oil. The crude product is chromatographed over flash silica with hexane:ethyl acetate (1:1 to 100% EtOAc) to give 110.

[(2R)-Isobutyl-3-carbo-*tert*-butoxy-propanoic acid amide of 2-oxo-3-amino-*N*-(carbomethoxymethyl)-pyridinone (111). The acid **23** (2.5 g, 11 mmole) is dissolved in 15 mL of THF and cooled to -15°C. NMM (1.3 mL, 11 mmole) and isobutylchloroformate (11 mmole, 1.6 mL) are added. After stirring for 30 min., NMM (1.3 mL, 11 mmole) and **110** (2.0 g, 11 mmole) are added. The reaction mixture is stirred for 17 hr and then quenched with 1N HCl and extracted with EtOAc. The organic layer is washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered and evaporated to give an oil which is chromatographed over flash silica with EtOAc to give **111**.

[(2R)-Isobutyl-3-*O*-Benzylhydroxylcarboxamido]-propanoic acid amide of 2-oxo-3-amino-*N*-(carbomethoxymethyl)-pyridinone (113). The ester **111** (0.8 g, 2 mmole) is dissolved in 5 mL of methylene chloride and trifluoroacetic acid (5 mL) is added slowly at 0°C. The mixture is stirred for 2 hr and evaporated to give **112**. The crude acid **112** is mixed with *O*-benzylhydroxylamine hydrochloride (0.5 g, 3.0 mmole), HOBT (0.81 g, 6 mmole), EDAC (0.58 g, 3 mmole) in 10 mL of DMF and 0.77 mL (7 mmole) of NMM at 0°C, and stirred for 17 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered and evaporated to give an oil which is chromatographed over flash silica with hexane:EtOAc (2:3 to 1:4) to give **113**.

[(2R)-Isobutyl-3-(*N*-hydroxylcarboxamido)]-propanoic acid amide of 2-oxo-3-amino-*N*-(carbomethoxymethyl)-pyridinone (22). The benzyl hydroxamic acid **113** (180 mg, 0.4 mmole) is taken in 5 mL of EtOH and the mixture is charged with 10% palladium on carbon (20 mg) and stirred under one atmosphere of hydrogen for 6 hours. TLC (EtOAc) indicates persistence of a small amount of starting material left. More palladium on carbon (18 mg) is added, **113** is and hydrogenated for another 20 min. The mixture is then filtered through celite and concentrated to give a semisolid, which is purified by preparative TLC (silica, EtOAc) to give **22**.

Example A

A tablet composition for oral administration, according to the present invention, is made comprising:

<u>Component</u>	<u>Amount</u>
•(2R)-[(1S)- <i>N</i> -Hydroxycarboxamido]-ethyl decanoic acid amide of <i>N</i> -[(2-methoxy)ethyl]- caprolactam-(3S)-amine ¹	15. mg
•Lactose	120. mg
•Maize Starch	70. mg

[(2R)-Isobutyl-3-carbo-*tert*-butoxy-propanoic acid amide of 2-oxo-3-amino-*N*-(carbomethoxymethyl)-pyridinone (111). The acid 23 (2.5 g, 11 mmole) is dissolved in 15 mL of THF and cooled to -15°C. NMM (1.3 mL, 11 mmole) and isobutylchloroformate (11 mmole, 1.6 mL) are added. After stirring for 30 min., NMM (1.3 mL, 11 mmole) and 110 (2.0 g, 11 mmole) are added. The reaction mixture is stirred for 17 hr and then quenched with 1N HCl and extracted with EtOAc. The organic layer is washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered and evaporated to give an oil which is chromatographed over flash silica with EtOAc to give 111.

[(2R)-Isobutyl-3-*O*-Benzylhydroxylcarboxamido]-propanoic acid amide of 2-oxo-3-amino-*N*-(carbomethoxymethyl)-pyridinone (113). The ester 111 (0.8 g, 2 mmole) is dissolved in 5 mL of methylene chloride and trifluoroacetic acid (5 mL) is added slowly at 0°C. The mixture is stirred for 2 hr and evaporated to give 112. The crude acid 112 is mixed with *O*-benzylhydroxylamine hydrochloride (0.5 g, 3.0 mmole), HOBT (0.81 g, 6 mmole), EDAC (0.58 g, 3 mmole) in 10 mL of DMF and 0.77 mL (7 mmole) of NMM at 0°C, and stirred for 17 hr at room temperature. The reaction is then partitioned between water and EtOAc. The organic layer is then washed with 1N HCl, NaHCO₃, and brine, dried over MgSO₄, filtered and evaporated to give an oil which is chromatographed over flash silica with hexane:EtOAc (2:3 to 1:4) to give 113.

[(2R)-Isobutyl-3-(*N*-hydroxylcarboxamido)]-propanoic acid amide of 2-oxo-3-amino-*N*-(carbomethoxymethyl)-pyridinone (22). The benzyl hydroxamic acid 113 (180 mg, 0.4 mmole) is taken in 5 mL of EtOH and the mixture is charged with 10% palladium on carbon (20 mg) and stirred under one atmosphere of hydrogen for 6 hours. TLC (EtOAc) indicates persistence of a small amount of starting material left. More palladium on carbon (18 mg) is added, 113 is and hydrogenated for another 20 min. The mixture is then filtered through celite and concentrated to give a semisolid, which is purified by preparative TLC (silica, EtOAc) to give 22.

Example A

A tablet composition for oral administration, according to the present invention, is made comprising:

<u>Component</u>	<u>Amount</u>
•(2R)-[(1S)- <i>N</i> -Hydroxycarboxamido]-ethyl decanoic acid amide of <i>N</i> -[(2-methoxy)ethyl]- caprolactam-(3S)-amine ¹	15. mg
•Lactose	120. mg
•Maize Starch	70. mg

- Talc 4. mg
- Magnesium Stearate 1. mg

1: The hydroxamic acid prepared according to Example 19. Other compounds having a structure according to Formula (I) are used with substantially similar results.

Example B

A capsule for oral administration, according to the present invention, is made comprising:

<u>Component</u>	<u>Amount (%w/w)</u>
• (2R)-[(1S)-N-hydroxycarboxamido]-ethyldecanoic acid amide of 1N-(carbomethoxy-methyl)-caprolactam-(3S)-amine ²	15%
• Polyethylene glycol	85%

2: The hydroxamic acid prepared according to Example 17. Other compounds having a structure according to Formula (I) are used with substantially similar results.

Example C

A saline-based composition for local administration, according to the present invention, is made comprising:

<u>Component</u>	<u>Amount (%w/w)</u>
• (2R)-Isobutyl-(3S)-[N-hydroxycarboxamido]-6-hydroxyhexanoic acid amide of (1N)-carbomethoxymethyl-caprolactam-(3S)-amine ³	10%
• Ethanol	10%
• Saline	80%

3: The hydroxamic acid prepared according to Example 16. Other compounds having a structure according to Formula (I) are used with substantially similar results.

- Talc 4. mg
- Magnesium Stearate 1. mg

1: The hydroxamic acid prepared according to Example 19. Other compounds having a structure according to Formula (I) are used with substantially similar results.

Example B

A capsule for oral administration, according to the present invention, is made comprising:

<u>Component</u>	<u>Amount (%w/w)</u>
•(2R)-[(1S)-N-hydroxycarboxamido]-ethyldecanoic acid amide of 1N-(carbomethoxy-methyl)-caprolactam-(3S)-amine ²	15%
•Polyethylene glycol	85%

2: The hydroxamic acid prepared according to Example 17. Other compounds having a structure according to Formula (I) are used with substantially similar results.

Example C

A saline-based composition for local administration, according to the present invention, is made comprising:

<u>Component</u>	<u>Amount (%w/w)</u>
•(2R)-Isobutyl-(3S)-[N-hydroxycarboxamido]-6-hydroxyhexanoic acid amide of (1N)-carbomethoxymethyl-caprolactam-(3S)-amine ³	10%
•Ethanol	10%
•Saline	80%

3: The hydroxamic acid prepared according to Example 16. Other compounds having a structure according to Formula (I) are used with substantially similar results.

Example D

A human female subject weighing 60 kg (132 lbs), suffering from rheumatoid arthritis, is treated by a method of this invention. Specifically, for 2 years, a tablet containing 50 mg of (2R)-*N*-hydroxycarboxamidemethyldecanoic acid amide of 1*N*-(carbomethoxy-methyl)-caprolactam-(3S)-amine (prepared according to Example 9) is administered orally to said subject.

At the end of the treatment period, the patient is examined and is found to have reduced inflammation, and improved mobility without concomitant pain.

Other compounds having a structure according to Formula (I) are used with substantially similar results.

Example E

A human male subject weighing 90 kg (198 lbs), suffering from osteoarthritis, is treated by a method of this invention. Specifically, for 5 years, a capsule containing 70 mg of (2R)-Isobutyl-3-(*N*-hydroxycarboxamido)-propanoic acid amide of *N*-(carbomethoxymethyl)-valerolactam-(3S)-amine (made according to Example 21) is administered daily to said subject.

At the end of the treatment period, the patient is examined via orthoscopy, and found to have no further advancement of erosion/fibrillation of the articular cartilage.

Other compounds having a structure according to Formula I are used with substantially similar results.

Example F

A human male subject weighing 90 kg (198 lbs), suffering from corneal ulcerations, is treated by a method of this invention. Specifically, for 2 months, a saline solution containing 10 mg of (2R)-*N*-Hydroxycarboxamidomethyl heptanoic acid amide of *N*-(carbomethoxymethyl)-caprolactam-(3S)-amine (made according to Example 8) is administered to said subject's affected eye twice-daily.

Other compounds having a structure according to Formula I are used with substantially similar results.

Example D

A human female subject weighing 60 kg (132 lbs), suffering from rheumatoid arthritis, is treated by a method of this invention. Specifically, for 2 years, a tablet containing 50 mg of (2R)-*N*-hydroxycarboxamidemethyldecanoic acid amide of 1*N*-(carbomethoxy-methyl)-caprolactam-(3S)-amine (prepared according to Example 9) is administered orally to said subject.

At the end of the treatment period, the patient is examined and is found to have reduced inflammation, and improved mobility without concomitant pain.

Other compounds having a structure according to Formula (I) are used with substantially similar results.

Example E

A human male subject weighing 90 kg (198 lbs), suffering from osteoarthritis, is treated by a method of this invention. Specifically, for 5 years, a capsule containing 70 mg of (2R)-Isobutyl-3-(*N*-hydroxycarboxamido)-propanoic acid amide of *N*-(carbomethoxymethyl)-valerolactam-(3S)-amine (made according to Example 21) is administered daily to said subject.

At the end of the treatment period, the patient is examined via orthoscopy, and found to have no further advancement of erosion/fibrillation of the articular cartilage.

Other compounds having a structure according to Formula I are used with substantially similar results.


Example F

A human male subject weighing 90 kg (198 lbs), suffering from corneal ulcerations, is treated by a method of this invention. Specifically, for 2 months, a saline solution containing 10 mg of (2R)-*N*-Hydroxycarboxamidomethyl heptanoic acid amide of *N*-(carbomethoxymethyl)-caprolactam-(3S)-amine (made according to Example 8) is administered to said subject's affected eye twice-daily.

Other compounds having a structure according to Formula I are used with substantially similar results.

What is Claimed is:

1. A compound having a structure according to Formula I




wherein

- (A) (1) (a) R^1 is hydrogen; alkyl; heteroalkyl; alkenyl; a heterocyclic ring; a carbocyclic ring; alkoxy; carbocycle-alkyl; heterocycle-alkyl; carbocycle-heteroalkyl; or heterocycle-heteroalkyl; and
- (b) R^2 is hydrogen; hydroxy; alkyl; alkenyl; alkynyl; heteroalkyl; a heterocyclic ring; a carbocyclic ring; carbocycle-alkyl; heterocycle-alkyl; or -OR, where R is alkyl, alkenyl, or carbocycle-alkyl; or
- (2) R^1 and R^2 together form a cycloalkyl ring having from 3 to 8 ring atoms;
- (B) R^3 is hydrogen; alkyl; or carbocycle-alkyl;
- (C) R^4 is
- (1) alkyl;
- (2) carbocycle-alkyl;
- (3) -X-C(=Y)-Z- R^5 or -X-CH₂-Z- R^5 , where
- (a) X is covalent bond or alkyl;
- (b) Y is O, S, or NH;
- (c) Z is O, S, or NH; and
- (d) R^5 is hydrogen; alkyl; alkenyl; carbocycle-alkyl; or aryl; or
- (4) -SO₂- R^6 , where R^6 is alkyl, carbocycle-alkyl, heterocycle-alkyl, or aryl; and
- (D) Q is -[C(R^7)₂]-_n, where
- (1) n is the integer 2, 3, or 4; and
- (2) each R^7 is independently hydrogen or alkyl so the Q-containing heterocycle is saturated; or the R^7 moiety on two adjacent carbon atoms is a covalent bond such that the Q-containing heterocycle in Formula (I) is unsaturated;

What is Claimed is:

1. A compound having a structure according to Formula I




wherein

- (A) (1) (a) R^1 is hydrogen; alkyl; heteroalkyl; alkenyl; a heterocyclic ring; a carbocyclic ring; alkoxy; carbocycle-alkyl; heterocycle-alkyl; carbocycle-heteroalkyl; or heterocycle-heteroalkyl; and
- (b) R^2 is hydrogen; hydroxy; alkyl; alkenyl; alkynyl; heteroalkyl; a heterocyclic ring; a carbocyclic ring; carbocycle-alkyl; heterocycle-alkyl; or -OR, where R is alkyl, alkenyl, or carbocycle-alkyl; or
- (2) R^1 and R^2 together form a cycloalkyl ring having from 3 to 8 ring atoms;
- (B) R^3 is hydrogen; alkyl; or carbocycle-alkyl;
- (C) R^4 is
- (1) alkyl;
- (2) carbocycle-alkyl;
- (3) -X-C(=Y)-Z- R^5 or -X-CH₂-Z- R^5 , where
- (a) X is covalent bond or alkyl;
- (b) Y is O, S, or NH;
- (c) Z is O, S, or NH; and
- (d) R^5 is hydrogen; alkyl; alkenyl; carbocycle-alkyl; or aryl; or
- (4) -SO₂- R^6 , where R^6 is alkyl, carbocycle-alkyl, heterocycle-alkyl, or aryl; and
- (D) Q is -[C(R^7)₂]-_n, where
- (1) n is the integer 2, 3, or 4; and
- (2) each R^7 is independently hydrogen or alkyl so the Q-containing heterocycle is saturated; or the R^7 moiety on two adjacent carbon atoms is a covalent bond such that the Q-containing heterocycle in Formula (I) is unsaturated;

or a pharmaceutically-acceptable salt, or biohydrolyzable alkoxyamide, acyloxyamide, or imide thereof.

2. The compound of Claim 1, wherein R^3 is hydrogen.
3. The compound of Claim 2, wherein R^1 is hydrogen; alkyl; or heteroalkyl; preferably R^1 is hydrogen; methyl; n-propyl; or 3-hydroxypropyl.
4. The compound of Claim 2, wherein R^2 is hydrogen; hydroxy; alkyl; alkenyl; or -OR, where R is alkyl or alkenyl; preferably wherein R^2 is C_1 - C_8 alkyl or C_2 - C_8 alkenyl; more preferably wherein R^2 is n-pentyl; n-octyl; or 2-methylpropyl.
5. The compound of Claim 4, wherein R^4 is carbocycle-alkyl; -X-C(=Y)-Z- R^5 ; or -X-CH₂-Z- R^5 .
6. The compound of Claim 5, wherein:
 - (a) when R^4 is carbocycle-alkyl, the alkyl is C_1 - C_4 alkyl and the carbocycle is aryl;
 - (b) when R^4 is -X-C(=Y)-Z- R^5 , X is C_1 ; Y is O; Z is O; and R^5 is alkyl, carbocycle-alkyl, or aryl;
 - (c) when R^4 is -X-C(=Y)-Z- R^5 , X is C_1 ; Y is O; Z is NH; and R^5 is alkyl, carbocycle-alkyl, or aryl; or
 - (d) when R^4 is -X-CH₂-Z- R^5 , X is C_1 ; Z is O or S; and R^5 is C_1 - C_3 .
7. A compound having a structure according to Formula (I)




wherein

- (A) (i) R^1 is hydrogen; alkyl; or heteroalkyl; and
(ii) R^2 is alkyl;
- (B) R^3 is hydrogen;
- (C) R^4 is

or a pharmaceutically-acceptable salt, or biohydrolyzable alkoxyamide, acyloxyamide, or imide thereof.

2. The compound of Claim 1, wherein R^3 is hydrogen.
3. The compound of Claim 2, wherein R^1 is hydrogen; alkyl; or heteroalkyl; preferably R^1 is hydrogen; methyl; n-propyl; or 3-hydroxypropyl.
4. The compound of Claim 2, wherein R^2 is hydrogen; hydroxy; alkyl; alkenyl; or -OR, where R is alkyl or alkenyl; preferably wherein R^2 is C_1 - C_8 alkyl or C_2 - C_8 alkenyl; more preferably wherein R^2 is n-pentyl; n-octyl; or 2-methylpropyl.
5. The compound of Claim 4, wherein R^4 is carbocycle-alkyl; -X-C(=Y)-Z- R^5 ; or -X-CH₂-Z- R^5 .
6. The compound of Claim 5, wherein:
 - (a) when R^4 is carbocycle-alkyl, the alkyl is C_1 - C_4 alkyl and the carbocycle is aryl;
 - (b) when R^4 is -X-C(=Y)-Z- R^5 , X is C_1 ; Y is O; Z is O; and R^5 is alkyl, carbocycle-alkyl, or aryl;
 - (c) when R^4 is -X-C(=Y)-Z- R^5 , X is C_1 ; Y is O; Z is NH; and R^5 is alkyl, carbocycle-alkyl, or aryl; or
 - (d) when R^4 is -X-CH₂-Z- R^5 , X is C_1 ; Z is O or S; and R^5 is C_1 - C_3 .
7. A compound having a structure according to Formula (I)



wherein

- (A) (i) R^1 is hydrogen; alkyl; or heteroalkyl; and
(ii) R^2 is alkyl;
- (B) R^3 is hydrogen;
- (C) R^4 is

- (1) alkyl;
 - (2) carbocycle-alkyl;
 - (3) $-X-C(=Y)-Z-R^5$ or $-X-CH_2-Z-R^5$, where
 - (a) X is covalent bond or alkyl;
 - (b) Y is O;
 - (c) Z is O or NH; and
 - (d) R^5 is alkyl; carbocycle-alkyl; or aryl; or
 - (4) $-SO_2-R^6$, where R^6 is alkyl, carbocycle-alkyl, heterocycle-alkyl, or aryl; and
- (D) Q is $[-C(R^7)_2-]_n$, where n is 4 and each R^7 is independently hydrogen or alkyl so the Q-containing heterocycle is saturated; or n is 4 and the R^7 moiety on two adjacent carbon atoms is a covalent bond such that the Q-containing heterocycle in Formula (I) is unsaturated; or a pharmaceutically-acceptable salt, biohydrolyzable amide or biohydrolyzable ester thereof.

8. The compound of Claim 7, wherein R^4 is $-X-C(=Y)-Z-R^5$.

9. The compound of Claim 8, wherein X is C_1-C_2 alkyl; Y is O; Z is O; and R^5 is methyl, benzyl, or t-butyl.

10. The compound of Claim 8, wherein X is C_1-C_2 alkyl; Y is O; Z is NH; and R^5 is methyl, benzyl or n-butyl.

11. The compound of Claim 7, wherein R^4 is $-X-CH_2-Z-R^5$.

12. The compound of Claim 11, wherein X is covalent bond or C_1-C_2 ; Z is O; and R^5 is C_1-C_3 .

13. The compound of Claim 1, wherein the compound is selected from:
 (2R)-Isobutyl-3-(N-hydroxycarboxamido)-propanoic acid amide of 1N-(carbomethoxy-methyl)-caprolactam-(3S)-amine;
 (2R)-Isobutyl-3-(N-hydroxycarboxamido)-propionic acid amide of 1N-(methyl-carboxamidomethyl)-caprolactam-(3S)-amine;
 (2R)-Isobutyl-3-(N-hydroxycarboxamido)-propionic acid amide of 1N-(carbo-*tert*-butoxy-methyl)-caprolactam-(3S)-amine;

- (1) alkyl;
 - (2) carbocycle-alkyl;
 - (3) $-X-C(=Y)-Z-R^5$ or $-X-CH_2-Z-R^5$, where
 - (a) X is covalent bond or alkyl;
 - (b) Y is O;
 - (c) Z is O or NH; and
 - (d) R^5 is alkyl; carbocycle-alkyl; or aryl; or
 - (4) $-SO_2-R^6$, where R^6 is alkyl, carbocycle-alkyl, heterocycle-alkyl, or aryl; and
- (D) Q is $[-C(R^7)_2-]_n$, where n is 4 and each R^7 is independently hydrogen or alkyl so the Q-containing heterocycle is saturated; or n is 4 and the R^7 moiety on two adjacent carbon atoms is a covalent bond such that the Q-containing heterocycle in Formula (I) is unsaturated; or a pharmaceutically-acceptable salt, biohydrolyzable amide or biohydrolyzable ester thereof.

8. The compound of Claim 7, wherein R^4 is $-X-C(=Y)-Z-R^5$.

9. The compound of Claim 8, wherein X is C_1-C_2 alkyl; Y is O; Z is O; and R^5 is methyl, benzyl, or t-butyl.

10. The compound of Claim 8, wherein X is C_1-C_2 alkyl; Y is O; Z is NH; and R^5 is methyl, benzyl or n-butyl.

11. The compound of Claim 7, wherein R^4 is $-X-CH_2-Z-R^5$.

12. The compound of Claim 11, wherein X is covalent bond or C_1-C_2 ; Z is O; and R^5 is C_1-C_3 .

13. The compound of Claim 1, wherein the compound is selected from:
 (2R)-Isobutyl-3-(N-hydroxycarboxamido)-propanoic acid amide of 1N-(carbomethoxy-methyl)-caprolactam-(3S)-amine;
 (2R)-Isobutyl-3-(N-hydroxycarboxamido)-propionic acid amide of 1N-(methyl-carboxamidomethyl)-caprolactam-(3S)-amine;
 (2R)-Isobutyl-3-(N-hydroxycarboxamido)-propionic acid amide of 1N-(carbo-*tert*-butoxy-methyl)-caprolactam-(3S)-amine;

(2R)-Isobutyl-3-(*N*-hydroxycarboxamido)-propionic acid amide of 1*N*-Benzyl-caprolactam-(3S)-amine;

(2R)-Isobutyl-3-*N*-hydroxycarboxamidopropionic acid amide of 1*N*-(carbo-benzyloxy-methyl)-caprolactam-(3S)-amine;

(2R)-Isobutyl-3-(*N*-hydroxycarboxamido)-propionic acid amide of 1*N*-(*N*-benzylcarboxamidomethyl)-caprolactam-(3S)-amine;

(2R)-Isobutyl-3-*N*-hydroxycarboxamidopropionic acid amide of 1*N*-(*n*-butyl-carboxamidomethyl)-caprolactam-(3S)-amine;

(2R)-*N*-Hydroxycarboxamidomethyl heptanoic acid amide of *N*-(carbomethoxymethyl)-caprolactam-(3S)-amine;

(2R)-*N*-hydroxycarboxamidomethyldecanoic acid amide of 1*N*-(carbomethoxymethyl)-caprolactam-(3S)-amine;

(2R)-*N*-Hydroxycarboxamidomethyl decanoic acid amide of *N*-toluenesulfonyl-caprolactam-(3S)-amine;

(2R)-*N*-Hydroxycarboxamidomethyl decanoic acid amide of *N*-[(2-methoxy)ethyl]-caprolactam-(3S)-amine;

(2R)-*N*-Hydroxycarboxamidomethyl decanoic acid amide of 1-*N*-*n*-butyl-caprolactam-(3S)-amine;

(2R)-Isobutyl-(3S)-[*N*-hydroxycarboxamido]-butanoic acid amide of (1*N*)-carbomethoxymethyl-caprolactam-(3S)-amine;

(2R)-Isobutyl-(3R)-[*N*-hydroxycarboxamido]-butanoic acid amide of (1*N*)-carbomethoxymethyl-caprolactam-(3S)-amine;

(2R)-Isobutyl-(3S)-[*N*-hydroxycarboxamido]-hexanoic acid amide of (1*N*)-carbomethoxymethyl-caprolactam-(3S)-amine;

(2R)-Isobutyl-(3S)-[*N*-hydroxycarboxamido]-6-hydroxyhexanoic acid amide of (1*N*)-carbomethoxymethyl-caprolactam-(3S)-amine;

(2R)-[(1S)-*N*-hydroxycarboxamido]-ethyldecanoic acid amide of 1*N*-(carbomethoxy-methyl)-caprolactam-(3S)-amine;

(2R)-[(1R)-*N*-hydroxycarboxamido]-ethyldecanoic acid amide of 1*N*-(carbomethoxy-methyl)-caprolactam-(3S)-amine;

(2R)-[(1S)-*N*-Hydroxycarboxamido]-ethyl decanoic acid amide of *N*-[(2-methoxy)ethyl]-caprolactam-(3S)-amine;

(2R)-[(1R)-*N*-Hydroxycarboxamido]-ethyl decanoic acid amide of *N*-[(2-methoxy)ethyl]-caprolactam-(3S)-amine;

(2R)-Isobutyl-3-(*N*-hydroxycarboxamido)-propanoic acid amide of *N*-(carbomethoxymethyl)-valerolactam-(3S)-amine; and

(2R)-Isobutyl-3-(*N*-hydroxycarboxamido)-propionic acid amide of 1*N*-Benzyl-caprolactam-(3S)-amine;

(2R)-Isobutyl-3-*N*-hydroxycarboxamidopropionic acid amide of 1*N*-(carbo-benzyloxy-methyl)-caprolactam-(3S)-amine;

(2R)-Isobutyl-3-(*N*-hydroxycarboxamido)-propionic acid amide of 1*N*-(*N*-benzylcarboxamidomethyl)-caprolactam-(3S)-amine;

(2R)-Isobutyl-3-*N*-hydroxycarboxamidopropionic acid amide of 1*N*-(*n*-butyl-carboxamidomethyl)-caprolactam-(3S)-amine;

(2R)-*N*-Hydroxycarboxamidomethyl heptanoic acid amide of *N*-(carbomethoxymethyl)-caprolactam-(3S)-amine;

(2R)-*N*-hydroxycarboxamidomethyldecanoic acid amide of 1*N*-(carbomethoxymethyl)-caprolactam-(3S)-amine;

(2R)-*N*-Hydroxycarboxamidomethyl decanoic acid amide of *N*-toluenesulfonyl-caprolactam-(3S)-amine;

(2R)-*N*-Hydroxycarboxamidomethyl decanoic acid amide of *N*-[(2-methoxy)ethyl]-caprolactam-(3S)-amine;

(2R)-*N*-Hydroxycarboxamidomethyl decanoic acid amide of 1-*N*-*n*-butyl-caprolactam-(3S)-amine;

(2R)-Isobutyl-(3S)-[*N*-hydroxycarboxamido]-butanoic acid amide of (1*N*)-carbomethoxymethyl-caprolactam-(3S)-amine;

(2R)-Isobutyl-(3R)-[*N*-hydroxycarboxamido]-butanoic acid amide of (1*N*)-carbomethoxymethyl-caprolactam-(3S)-amine;

(2R)-Isobutyl-(3S)-[*N*-hydroxycarboxamido]-hexanoic acid amide of (1*N*)-carbomethoxymethyl-caprolactam-(3S)-amine;

(2R)-Isobutyl-(3S)-[*N*-hydroxycarboxamido]-6-hydroxyhexanoic acid amide of (1*N*)-carbomethoxymethyl-caprolactam-(3S)-amine;

(2R)-[(1S)-*N*-hydroxycarboxamido]-ethyldecanoic acid amide of 1*N*-(carbomethoxy-methyl)-caprolactam-(3S)-amine;

(2R)-[(1R)-*N*-hydroxycarboxamido]-ethyldecanoic acid amide of 1*N*-(carbomethoxy-methyl)-caprolactam-(3S)-amine;

(2R)-[(1S)-*N*-Hydroxycarboxamido]-ethyl decanoic acid amide of *N*-[(2-methoxy)ethyl]-caprolactam-(3S)-amine;

(2R)-[(1R)-*N*-Hydroxycarboxamido]-ethyl decanoic acid amide of *N*-[(2-methoxy)ethyl]-caprolactam-(3S)-amine;

(2R)-Isobutyl-3-(*N*-hydroxycarboxamido)-propanoic acid amide of *N*-(carbomethoxymethyl)-valerolactam-(3S)-amine; and

[(2R)-Isobutyl-3-(*N*-hydroxylcarboxamido)]-propanoic acid amide of 2-oxo-3-amino-*N*-(carbomethoxymethyl)-pyridinone.

14. A pharmaceutical composition comprising:
- (a) a safe and effective amount of a compound of Claim 1 to Claim 13; and
 - (b) a pharmaceutically-acceptable carrier.

15. A method for preventing or treating a disease associated with unwanted matrixmetalloprotease activity in a human or other animal subject, the method comprising administering to said subject a safe and effective amount of a compound of Claim 1 to Claim 13.

[(2R)-Isobutyl-3-(*N*-hydroxylcarboxamido)]-propanoic acid amide of 2-oxo-3-amino-*N*-(carbomethoxymethyl)-pyridinone.

14. A pharmaceutical composition comprising:
- (a) a safe and effective amount of a compound of Claim 1 to Claim 13; and
 - (b) a pharmaceutically-acceptable carrier.

15. A method for preventing or treating a disease associated with unwanted matrixmetalloprotease activity in a human or other animal subject, the method comprising administering to said subject a safe and effective amount of a compound of Claim 1 to Claim 13.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 96/03726

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D223/12 C07D211/76 C07D213/75 A61K31/55 A61K31/44
A61K31/44

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP,A,0 236 872 (F. HOFFMANN - LA ROCHE AG) 16 September 1987 cited in the application see the whole document ---	1-14
Y	WO,A,90 05716 (BRITISH BIO-TECHNOLOGY LIMITED) 31 May 1990 see the whole document ---	1-14
Y	WO,A,92 17460 (SMITHKLINE BEECHAM PLC) 15 October 1992 cited in the application see the whole document ---	1-14
Y	WO,A,94 21612 (OTSUKA PHARMACEUTICAL CO., LTD.) 29 September 1994 see the whole document ---	1-14
-/--		

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

A document member of the same patent family

Date of the actual completion of the international search

20 June 1996

Date of mailing of the international search report

07. 08. 96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+ 31-70) 340-3016

Authorized officer

Hartrampf, G

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 96/03726

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D223/12 C07D211/76 C07D213/75 A61K31/55 A61K31/44
A61K31/44

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP,A,0 236 872 (F. HOFFMANN - LA ROCHE AG) 16 September 1987 cited in the application see the whole document ---	1-14
Y	WO,A,90 05716 (BRITISH BIO-TECHNOLOGY LIMITED) 31 May 1990 see the whole document ---	1-14
Y	WO,A,92 17460 (SMITHKLINE BEECHAM PLC) 15 October 1992 cited in the application see the whole document ---	1-14
Y	WO,A,94 21612 (OTSUKA PHARMACEUTICAL CO., LTD.) 29 September 1994 see the whole document ---	1-14
-/--		

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

A document member of the same patent family

Date of the actual completion of the international search

20 June 1996

Date of mailing of the international search report

07. 08. 96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+ 31-70) 340-3016

Authorized officer

Hartrampf, G

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 96/03726

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>WO,A,93 16982 (JAMES BLACK FOUNDATION LIMITED) 2 September 1993 see the whole document -----</p>	1-14

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 96/03726

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>WO,A,93 16982 (JAMES BLACK FOUNDATION LIMITED) 2 September 1993 see the whole document -----</p>	1-14

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 96/ 03726

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claim 15 is directed to a method of treatment of (diagnostic method practised on) the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 96/ 03726

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claim 15 is directed to a method of treatment of (diagnostic method practised on) the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/US 96/03726

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-236872	16-09-87	AU-B- 588437	14-09-89
		AU-B- 6990287	17-09-87
		CA-A- 1314655	16-03-93
		DE-A- 3782751	07-01-93
		IE-B- 60128	01-06-94
		JP-B- 6029228	20-04-94
		JP-A- 62230757	09-10-87
		US-A- 4996358	26-02-91

WO-A-9005716	31-05-90	AU-B- 641629	30-09-93
		AU-B- 4746890	12-06-90
		CA-A- 2003719	23-05-90
		DE-D- 68913988	21-04-94
		DE-T- 68913988	28-07-94
		EP-A- 0445206	11-09-91
		ES-T- 2063334	01-01-95
		JP-T- 4503057	04-06-92
		NO-B- 177700	31-07-95
		US-A- 5514677	07-05-96
		US-A- 5304604	19-04-94

WO-A-9217460	15-10-92	EP-A- 0579650	26-01-94
		JP-T- 6505989	07-07-94

WO-A-9421612	29-09-94	AU-B- 6263794	11-10-94
		CA-A- 2136108	29-09-94
		CN-A- 1105799	26-07-95
		EP-A- 0641323	08-03-95
		JP-A- 7157470	20-06-95

WO-A-9316982	02-09-93	AU-B- 3509793	13-09-93
		CA-A- 2128998	02-09-93
		EP-A- 0626942	07-12-94
		FI-A- 943817	19-08-94
		HU-A- 71499	28-11-95
		JP-T- 7504184	11-05-95
		NO-A- 943055	11-10-94
		US-A- 5514683	07-05-96
		ZA-A- 9301193	19-08-94

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 96/03726

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-236872	16-09-87	AU-B- 588437	14-09-89
		AU-B- 6990287	17-09-87
		CA-A- 1314655	16-03-93
		DE-A- 3782751	07-01-93
		IE-B- 60128	01-06-94
		JP-B- 6029228	20-04-94
		JP-A- 62230757	09-10-87
		US-A- 4996358	26-02-91

WO-A-9005716	31-05-90	AU-B- 641629	30-09-93
		AU-B- 4746890	12-06-90
		CA-A- 2003719	23-05-90
		DE-D- 68913988	21-04-94
		DE-T- 68913988	28-07-94
		EP-A- 0445206	11-09-91
		ES-T- 2063334	01-01-95
		JP-T- 4503057	04-06-92
		NO-B- 177700	31-07-95
		US-A- 5514677	07-05-96
		US-A- 5304604	19-04-94

WO-A-9217460	15-10-92	EP-A- 0579650	26-01-94
		JP-T- 6505989	07-07-94

WO-A-9421612	29-09-94	AU-B- 6263794	11-10-94
		CA-A- 2136108	29-09-94
		CN-A- 1105799	26-07-95
		EP-A- 0641323	08-03-95
		JP-A- 7157470	20-06-95

WO-A-9316982	02-09-93	AU-B- 3509793	13-09-93
		CA-A- 2128998	02-09-93
		EP-A- 0626942	07-12-94
		FI-A- 943817	19-08-94
		HU-A- 71499	28-11-95
		JP-T- 7504184	11-05-95
		NO-A- 943055	11-10-94
		US-A- 5514683	07-05-96
		ZA-A- 9301193	19-08-94
